

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



7009000

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 48/00, C07K 14/47, C12N 1/15, 1/21, 5/10, 15/09, 15/12, C12P 21/02</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/31396</b>
			<b>(43) International Publication Date:</b> 23 July 1998 (23.07.98)
<b>(21) International Application Number:</b> PCT/US97/06864		<b>Quennet, F-78955 Carrières-sous-Poissy (FR). BOUILLAUD, Frederic [FR/FR]; 54, allée de la Forêt, F-92360 Meudon la Forêt (FR).</b>  <b>(74) Agent:</b> WILSON, Mary, J.; Nixon & Vanderhye P.C., 8th floor, 1100 North Glebe Road, Arlington, VA 22201-4714 (US).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
<b>(22) International Filing Date:</b> 22 April 1997 (22.04.97)			
<b>(30) Priority Data:</b> 60/034,960      15 January 1997 (15.01.97)      US			
<b>(71) Applicants (for all designated States except US):</b> DUKE UNIVERSITY [US/US]; 230 North Building, Research Drive, Box 90083, Durham, NC 27708-0083 (US). THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE CENTRE DE RECHERCHE SUR L'ENDOCRINOLOGIE MOLECULAIRE ET LE DEVELOPPEMENT [FR/FR]; 9, rue Jules Hetzel, F-92190 Muedon (FR).			
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SURWIT, Richard, S. [US/US]; 3804 Sweeten Creek Road, Chapel Hill, NC 27514 (US). COLLINS, Sheila, A. [US/US]; 1516 Woodburn Road, Durham, NC 27705 (US). WARDEN, Craig, H. [US/US]; 728 Waxwing Place, Davis, CA 95616 (US). SELDIN, Michael, F. [US/US]; 3207 Magpie, Davis, CA 95616 (US). RICQUIER, Daniel [FR/FR]; 167, rue Alexis		<b>Published</b> <i>With international search report.</i> <i>With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.</i> <i>Date of receipt by the International Bureau:</i> 23 February 1998 (23.02.1998)	
<b>(54) Title:</b> RESPIRATION UNCOUPLING PROTEIN			
<b>(57) Abstract</b>  The present invention relates, in general, to a protein linked to cell respiration, thermogenesis, obesity and hyperinsulinemia and, in particular, to a protein designated uncoupling protein-2 (UCP2) and to nucleic acid sequences encoding same. The invention also relates to diagnostic methodologies based, for example, on a determination of levels of UCP2 expression. Further, the invention relates to therapies involving modulating UCP2 expression and/or activity. In addition, the present invention relates to methods of screening compounds for their suitability for use in such therapies.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**RESPIRATION UNCOUPLING PROTEIN**

The entire content of Provisional Application No. 60/034,960, filed January 15, 1997, is incorporated herein by reference.

**TECHNICAL FIELD**

The present invention relates, in general, to a protein linked to cell respiration, metabolic rate, thermogenesis, obesity and hyperinsulinemia and, in particular, to a protein designated uncoupling protein-2 (UCP2) and to nucleic acid sequences encoding same. The invention also relates to diagnostic methodologies based, for example, on a determination of levels of UCP2 expression. Further, the invention relates to therapies involving modulating UCP2 expression and/or activity. In addition, the present invention relates to methods of screening compounds for their suitability for use in such therapies.

**BACKGROUND**

Obesity is a major health problem in most industrialized countries, including the United States. The seriousness of the pathologies associated with obesity, including type II diabetes and hypertension, has prompted numerous studies directed at determining the molecular basis for obesity. One result of such

studies is the demonstration that genetically inherited characteristics may play a significant role in the development of the disorder.

Recent reports indicate that obesity may be associated with an energy imbalance. The energy balance of an individual is dependent upon both the energy intake (food) and the energy output (corresponding to resting metabolic rate metabolism, physical exercise and heat production). One hypothesis is that decreased resting metabolic rate and/or thermogenesis contributes to the development of obesity. Results of studies relating to uncoupling protein-1 (UCP1) demonstrate that such is the case in rodents (Ricquier et al, FASEB J. 5:2237 (1991); Muzin et al, Biochem. J. 261:721 (1989); Ricquier et al, J. Biol. Chem. 261:13905 (1986); Collins et al, Endocrinol. 138:405 (1997)).

UCP1 creates a pathway that allows dissipation of the electrochemical gradient of protons across the inner mitochondrial membrane in brown adipose tissue, without coupling to any other energy consuming process. This results in generation of heat and dissipation of calories. This process has been implicated in the regulation of body temperature, body composition and glucose metabolism in animals. However, UCP1-containing brown adipose tissue is unlikely to play a role in weight regulation in adult animals and humans living in a thermoneutral

environment since there is little brown adipose tissue present.

The present invention relates to a novel uncoupling protein, designated UCP2. The chromosomal location of the UCP2 gene, and uncoupling activity and tissue distribution of UCP2 are consistent with a role for this protein in diabetes and obesity.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-F. A. Human UCP2 amino acid sequence. (Human UCP2 sequence is GenBank accession U76367.) The three mitochondrial carrier protein motifs present in UCP1 and conserved in UCP2 are underlined. B. Nucleotide sequence encoding human UCP2 amino acid sequence shown in A, including the stop codon. C. Map of the human UCP2 gene. D. Subcloned fragments of hUCP2-gl. pSU04, pSUB14sp6, pSUB 14t7, pSUB23 sp6, pSUB23t7 are non localized fragments of the human UCP2 gene. pSUB 12 SP6 corresponds to the 3' end of the gene; pSUB5t7 corresponds to the promoter region; ex 12 identifies a genomic region containing exon 1 and 2 plus intronic regions; ex 34 identifies a genomic region containing exon 3 and 4 plus intronic regions; and ex 56 identifies a genomic region containing exon 5 and 6 plus intronic regions. Cloning strategy: A mouse UCP cDNA was first cloned using rat UCP1 cDNA to screen a mouse muscle cDNA library (lambda GT11, Clontech).

Primers were derived from the mouse UCP2 cDNA to amplify a genomic human fragment corresponding to exons 1 and 2, plus introns 1 and 2 of the human UCP2 gene. This partial genomic fragment was then used to clone the entire human UCP2 gene, hUCP2-g1. The library screened was a human genomic library from Clontech (cat# HL 1067j; lot #45003) made in the EMBL3 SP6/T7 cloning vector with *E. coli* K802 as host strain. The DNA source was human placenta. E. Consensus sequence for the entire human expressed UCP2. The entire expressed sequence (mRNA) is 1612 bp long. The coding sequence extends from bp 345 to 1275 of this sequence. The clones hUCP2-5' and hUCP2-3', from which the consensus sequence was derived, were isolated from the human lung Marathon library from Clontech (Palo Alto, CA; catalog #7408-1). They were isolated by Random Amplification of cDNA Ends (RACE). 5' and 3' RACE were performed on the Marathon library using methods as described by Clontech. The 5' gene specific primer sequence was AGAGAAGGGAAGGAGGGAAG (hUCP2.CDSR3). The 3' gene specific primer sequence was CATCTCCTGGGACGTAGC (hUCP2.CDSF3). PCR conditions were 94°C 30 sec, 64°C 30 sec, 72°C 1 min for 30 cycles on an MJ Research PTC-200, with Clontech KlenTaq Plus DNA polymerase. F. Oligonucleotides that can be used to amplify the UCP2 coding sequence. Names are: hUCP2.CDS (F or R, pair number) where hUCP2 means human UCP2, CDS means coding sequence and F or R

refers to forward or reverse primer. PCR conditions are in an MJ research PTC-200 96-well machine. Clontech KlenTaq plus DNA polymerase were used: 94°C 30 sec, 60°C 30 sec, and 68°C 2 min. Times 30 cycles.

5 Figure 2. Flow cytometry analysis of membrane potential following the expression of UCP2 in *S. cerevisiae*; comparison of wild-type UCP1 and non-active UCP1 mutants. The number of cells (counts, Y-axis) is represented on a logarithmic scale of 1024  
10 channels. The X-axis is a logarithmic scale of fluorescence intensity. A shift of the curve towards the left indicates a decreased mitochondrial potential. The horizontal bar on the left indicates the position of cells treated with mCICCP, a chemical  
15 uncoupler. The expression of UCP2, UCP1 or UCPmut mRNA in recombinant yeast, as well as its absence in wild type yeast, was checked by Northern blots.

20 Figures 3A-C. A. Tissue distributions of UCP2 in humans and mice. Human UCP2 mRNA was detected on a human multiple tissue Northern blot containing 2  $\mu$ g mRNA per lane (Clontech, Palo Alto). This blot was probed with  $^{32}$ P-labeled human UCP2 insert from IMAGE clone 129216. The  $^{32}$ P-labeled insert was used to probe the Master blot in ExpressHyb solution (Clontech, Palo  
25 Alto, CA). Expression of UCP1 and UCP2 in mice was examined in various tissues of mice maintained at 23°C. 20  $\mu$ g RNA was loaded in each lane except the WAT lane which has 4  $\mu$ g. UCP1 mRNA size is 1.5 kb,

SUBSTITUTE SHEET (RULE 26)

while UCP2 mRNA size is 1.6 kb. The UCP1 probe is the HSU 28480 clone, while UCP2 clone is GenBank accession number U69135 (DR). Abbreviations: H, heart; B, brain; Pl, placenta; Lu, lung; L, liver; M, skeletal muscle; K, kidney; P, pancreas; Wat, white adipose tissue; Bat, brown adipose tissue; and M1 and M2 are thigh and abdominal muscle, respectively.

B. Chromosomal location of Ucp2 on distal mouse chromosome 7. The approximate position in centi-Morgan (cM) distance from the centromere, relationship among other genes and corresponding human chromosomal positions of orthologous genes are shown. The Ucp2 gene segregated in all 114 meiotic events examined with  $\beta$ -arrestin (*Arrb1*), a gene previously mapped to this position on both mouse chromosome 7 and to the long arm of human chromosome 11 (band q13). The other symbols and corresponding names are: *Tyr*, tyrosinase; *D7Was12*, DNA segment Chr 7, Washington 12; *Omp*, olfactory marker protein; *Hbb*, hemoglobin beta chain complex; *tub*, tubby mutation; *Calc*, calcitonin; and *D7Mit40*, DNA segment Chr 7 Mit 40.

C. Regulation of UCP2 mRNA levels in A/J and C57BL/6J by diet. 30  $\mu$ g of epididymal fat pad white adipose tissue RNA from 2 mice from each strain and diet was electrophoresed and a Northern blot prepared. The high-fat diet is the same as described previously (Surwit et al, Diabetes 37:1163 (1988)). The blot was probed with 32P-labeled clone 129216 (human UCP2) and



with cyclophilin. Amounts of UCP2 mRNA were determined with a Molecular Dynamics phosphorimager and were normalized with cyclophilin mRNA levels.

Figure 4. Effect of diet on UCP2 mRNA levels in A/J and B6 mice. Methods are as described in Figure 3C. Epididymal white adipose tissue was prepared from mice on a low fat diet (L), a high fat diet (Surwit et al, Diabetes 37:1163 (1988)) (H), or the high fat diet and the  $\beta 3$  adrenergic agonist CL316,243 (H $\beta$ ) for 25 days (Collins et al, Endocrin. 138:405 (1997)).

Figure 5. Hypothesis about the regulation of resting metabolic rate and thermogenesis by UCP1 and UCP2. In mitochondria, the proton electrochemical gradient ( $\Delta\mu H^+$ ) generated by the respiratory chain allows ATP synthesis by ATP synthase. In brown adipocyte mitochondria, uniquely, UCP1 short-circuits the proton circuit and oxidative energy is dissipated as heat in response to cold exposure. It has also been proposed that cafeteria diets may activate brown adipocyte thermogenesis (Rothwell et al, Nature 281:31 (1979)). In contrast, mitochondria of many tissues express UCP2, an homolog of UCP1, that may also lower the level of coupling of respiration to ADP phosphorylation. This may also lead to increased resting metabolic rate and body temperature and underlies "thermogenesis". The present data show that UCP2 is not induced by cold but is induced by diet.

Figure 6. Increase in UCP2 expression by thiazolidinediones.

Figure 7. Map showing organization of the mouse UCP2 gene.

5      Figures 8A and B. Mouse UCP2 sequences.

A. Partial promoter sequence. B. Promoter and coding sequence.

Figure 9. Map showing organization of the human UCP2 gene.

10      Figures 10A-D. Sequences of four regions of the human UCP2 gene.

Figure 11. Functional activity of UCP2 promoter. The following CAT reporter constructs were transfected into HIB-1B cells: SV2 contains the SV40 T-antigen promoter; SV0 is a promoter-less control construct; UCP2(+) and UCP2(-) indicates constructs containing the 246 bp fragment from the mouse UCP2 gene inserted in the "sense" orientation or the "antisense" orientation, respectively. Twenty-four hours following their introduction into HIB-1B, the cells were harvested, a soluble extract of the cells was prepared and the level of CAT activity was measured by the TLC method. The data shown are the mean  $\pm$  sd of two determinations, and the CAT activity is determined as acetylated chloramphenicol product/total, and is expressed as an arbitrary unit. The sample marked as Blank is the unreacted chloramphenicol substrate.

## 5

10

15

Objects and advantages of the present invention will be clear from the description that follows.

## 20

## 25

The present invention relates to a protein that is involved in energy balance, body weight regulation and thermoregulation. The protein, UCP2, is widely expressed in mammalian tissues, including human tissues, and UCP2 mRNA levels in white fat are elevated in response to fat feeding. The UCP2 gene sequence maps to a chromosomal region linked to obesity and hyperinsulinemia.

The identification of UCP2 provides an explanation for a proton leak that has been reported in mitochondria (Brand et al, Biochem. J. 275:81 (1991); Porter et al, Am. J. Physiol. 269:R1213 (1995); Porter et al, Nature 362:628 (1993)). This leak is related to the standard metabolic rate which varies inversely with body mass of various species and with hormonal status (Brand et al, Bio. Chem. J. 275:81 (1991); Porter et al, Nature 362:628 (1993); Rice et al, Obesity Research 4:441 (1996)). Thus, UCP2 proton conductants can be expected to vary with membrane potential, as is the case with UCP1 (Diolez et al, in 8th European Bioenergetic Conference, Valencia, Spain (1994)). This property permits the leak to operate without collapsing membrane potential.

In brown adipose tissue, UCP1 creates a pathway that allows dissipation of the electrochemical gradient of protons across the inner membrane, without coupling to any other energy consuming process (Nedergard et al, in Molecular Mechanisms and Bioenergetics 23:385-420, ed. Ernster, L., Elsevier Science (1992)). The consequences are lower efficiency of oxidative phosphorylation and increased heat production. Previously, this was expected to be a general property of the membrane (involving the phospholipid composition) or a side effect of a known protein such as the ADP/ATP carrier. As shown in Fig. 5, it is now known that UCP2, a thermogenic and

5

25

relates to methods of screening compounds for the ability to bind to or alter the activity of or the expression of the UCP2 gene product. In addition, the invention relates to diagnostic and treatment methodologies based on UCP2 and its encoding sequence.

UCP2 Encoding Sequence and Methods of UCP2 Production Using Same, UCP2 and Anti-UCP2 Antibodies

The present invention relates to nucleotide sequences that encode a protein that partially uncouples respiration (as does UCP1) in non brown (as well as brown) adipose tissue, for example, mammalian UCP2, particularly, human UCP2, or portions thereof as defined above (examples of such portions include sequences encoding the 10 N-terminal amino acids, and sequences encoding the mitochondrial carrier protein motifs of Fig. 1A). In particular, the present invention relates to nucleotide sequences that encode the amino acid sequence given in Figure 1A, or portions thereof as defined above (the specific DNA sequence encoding UCP2 given in Figure 1B being only an example). Further, nucleotide sequences to which the invention relates include those encoding substantially the same protein as shown in Figure 1A, for example, inter- and intra-species variations thereof (see Example IX), as well as functional equivalents of the amino acid sequence shown in Figure 1A. The invention further relates to nucleotide

sequences substantially identical to the sequence shown in Figure 1B. A "substantially identical" sequence is one the complement of which hybridizes to the nucleic acid sequence of Figure 1B at 42°C in 50% formamide; 1 X saline/sodium citrate (SSC) containing 0.25% SDS and which remains bound when subjected to washing at 52-55°C with 0.1-0.2 X SSC containing 0.1% SDS (note: 20 X SSC = 3 M sodium chloride/0.3 M sodium citrate). The invention also relates to nucleic acids complementary to those described above.

The present invention also relates to a recombinant molecule comprising a nucleotide sequence as described above and to a host cell transformed therewith. Using standard methodologies, well known in the art, a recombinant molecule comprising a vector and a nucleotide sequence encoding the UCP2 protein, or portion thereof as defined above, can be constructed. Vectors suitable for use in the present invention include plasmid and viral vectors. Plasmid vectors into which a nucleic sequence encoding the UCP2 protein, or portion thereof, can be cloned include any vectors compatible with transformation into a selected host cell. Such vectors include vectors suitable for introduction into yeast and insect cells, generally, mammalian expression vectors suitable for expression in host cells, which vectors can include sequence elements that enhance transcription and/or prolong mRNA halflife in the cell

(e.g.  $\beta$ -globin gene 3' untranslated region) specifically, pUC-based *E. coli* vectors, pYeDPUCP2, pSelectUCP2, and PECE-UCP2. The nucleotide sequence of the invention can be present in the vector operably  
5 linked to regulatory elements, for example, a promoter. Suitable promoters include, but are not limited to, tissue specific promoters (e.g. leptin gene promoter or aP2 gene promoter specific for  
10 adipose cells, muscle creatine kinase promoter specific for skeletal muscle and lymphoid cell promoters), muscle actin promoter, interleukin promoter, CMV, SV40 and MMTV promoters.

As indicated above, the recombinant molecule of the invention can be constructed so as to be suitable  
15 for transforming a host cell. Suitable host cells include prokaryotic cells, such as bacteria, lower eukaryotic cells, such as yeast, and higher eukaryotic cells, such as mammalian cells, and insect cells. The recombinant molecule of the invention can be  
20 introduced into appropriate host cells by one skilled in the art using a variety of known methods.

The present invention further relates to a method of producing UCP2, or portions thereof as defined  
25 above. The method comprises culturing the above-described transformed host cells under conditions such that the encoding sequence is expressed and the protein thereby produced.



In addition to the nucleic acids described above, the present invention also relates to UCP2 gene sequences, including introns, exons and flanking regions (e.g. the UCP2 promoter), and to portions thereof suitable for use as probes or primers. The invention also relates to nucleic acid sequences corresponding to the entire expressed UCP2 sequence (e.g. UCP2 mRNA or corresponding cDNA), as well as portions thereof suitable, for example, for use as probes or primers.

A human UCP2 genomic clone, hUCP2-g1, was deposited on January 13, 1997, and was given Accession No. I-1806. A further human genomic clone (designated hUCP2-g2 and a mouse genomic clone (designated MMU2-L2) were deposited at the Collection Nationale de Cultures de Microorganismes, Institut Pasteur, 25, Rue du Docteur Roux, 75724, Paris CEDEX 15, France, under the terms of the Budapest Treaty, on April 16, 1997, and were given Accession Nos. I-1867 and I-1868, respectively. hUCP2-g2 was cloned from the same library used to clone hUCP2-g1. A 500 bp DNA corresponding to the 5' end of hUCP2-g1 was used to screen the genomic library. MMU2-L2 was cloned from a mouse genomic library screened using the mouse UCP2 cDNA. The genomic library was from Strategene (La Jolla, C: 129SVJ Mouse genomic library -- catalog number 946306 -- in the Lamda FIX II vector; the amplification host was XL1-blue MRA(P2). Sequence 2

of hUCP2-g2 is 1161 bp long and goes from position  
-511 to + 650. Therefore, this fragment contains the  
putative human promoter region. A map of the human  
UCP2 gene is shown in Fig. 1C and the sequences of  
several subcloned fragments are given in Fig. 1D.  
(See also Figures 9 and 10.) Two further deposits  
(hUCP2-5' and hUCP2-3') were made at the American Type  
Culture Collection, 12301 Parklawn Drive, Rockville,  
MD 20852, USA, under the terms of the Budapest Treaty,  
on January 10, 1997. These deposits were given  
Accession Nos. 97850 and 97849, respectively. These  
deposits comprise plasmids containing inserts from  
normal human lung cDNA. The plasmids are in the  
Invitrogen pCR2.1 vector. hUCP2-5' includes sequence  
from base pair (bp) 1 up to bp 1375, thus including  
the entire 5' untranslated sequence and the entire  
coding sequence. hUCP2-3' includes sequence from bp  
313 up to bp 1612, thus including the entire coding  
sequence and the entire 3' untranslated region. A  
bacterial artificial chromosome (BAC) clone for human  
genomic UCP2 DNA has been isolated. This clone,  
hUCP2.BAC, was deposited with the American Type  
Culture Collection, 12301 Parklawn Drive, Rockville,  
MD 20852, USA, under the terms of the Budapest Treaty  
on April 18, 1997. The clone was isolated by  
hybridizing hUCP2 probe to the human BAC library  
prepared by Genome Systems. The probe was PCR product  
for human UCP2 produced from human lung cDNA and

amplified with hUCP2.cds3 primers as described in Example IX. BAC clone inserts range from 90 to 300 kb in length, with an average of 120 kb. Hybridization and isolation of hUCP2.BAC were performed by Genome Systems using their standard techniques. Since the human UCP2 gene is approximately 20 kb, hUCP2.BAC is believed to contain the entire gene as well as the entire promoter. hUCP2.BAC has also been digested with EcoRI and subcloned into the pZERO vector of Invitrogen. These subclones have been hybridized to hUCP2-5' probe to identify clones with the human promoter. They have also been hybridized to a (CA)<sub>14</sub> oligo to identify a polymorphic CA repeat. Consensus sequence for the entire human expressed UCP2 is shown in Fig. 1E.

Nucleic acid sequence(s) of the invention can be used, in accordance with standard protocols, as probes and primers. Oligonucleotides suitable for amplifying the human UCP2 coding sequence are given in Fig. 1F (pairs 1 and 3 exhibit little non-specific amplification).

The present invention further relates to mammalian UCP2, particularly, human UCP2, substantially free of proteins with which it is normally associated, or portions thereof as defined above. The proteins, polypeptides and peptides of the invention can be produced recombinantly using the nucleic acid sequences as described above, or

5

15

## 20

25

ability either to increase or to decrease expression or activity of UCP2. The assays are performed *in vitro* or *in vivo*. *In vitro*, cells expressing UCP2 are incubated in the presence and absence of the test compound. By determining the level of UCP2 expression in the presence of the test compound (using, for example, Northern blots, immunoassays (e.g. RIA, Western blots or immunohistochemistry) or PCR), or the level of UCP2 activity in the presence of the test compound, compounds can be identified that suppress or enhance UCP2 expression or activity. Alternatively, constructs comprising the UCP2 promoter operably linked to a reporter gene (e.g. luciferase, chloramphenicol acetyl transferase, LacZ, green fluorescent protein, etc.) can be introduced into host cells and the effect of the test compounds on expression of the reporter gene detected.

Cells suitable for use in the foregoing assays include, but are not limited to, lymphoblasts, myocytes, adipocytes and hepatic cells, more specifically, C2C12 cells, 3T3 cells of adipocyte lineage, HIB-1B cells, rodent hepatoma cells, HepG2 cells, and B7 cells. (See Example V.)

Compounds that suppress or enhance UCP2 expression can also be identified using *in vivo* screens. In these assays, the test compound is administered (e.g. IV, IP, IM, orally, or otherwise), to the animal, for example, at a variety of dose

5 levels. The effect of the compound on UCP2 expression is determined by comparing UCP2 levels, for example, in blood, muscle or fat tissue, using Northern blots, immunoassays, PCR, etc., as described above. Suitable  
10 test animals include rodents, primates, dogs and swine. Humanized mice can also be used as test animals, that is mice in which the endogenous mouse protein is ablated (knocked out) and the homologous human protein added back by standard transgenic  
15 approaches. Such mice express only the human form of a protein. Humanized mice expressing just the human UCP2 can be used to study *in vivo* responses of weight loss, fever, cachexia in response to potential agents regulating UCP2 protein or mRNA levels. As an  
20 example, transgenic mice have been produced carrying the human apoE4 gene. They are then bred with a mouse line that lacks endogenous apoE, to produce an animal model carrying human proteins believed to be instrumental in development of Alzheimers pathology. Such transgenic animals are useful for dissecting the  
25 biochemical and physiological steps of disease, and for development of therapies for disease intervention (Loring, et al, Neurobiol. Aging 17:173 (1996)).

Compounds that suppress or enhance UCP2 activity can be identified by contacting UCP2 with the test compound under conditions such that the compound can interact with (e.g. bind to) the protein. A system such as the yeast expression system described in

Example I can be used. In such a system, the effect of the test compound on UCP2 activity can be determined, for example, by analyzing the alteration in membrane potential (e.g. using flow cytometry) (see Example I). (Comparable studies can be carried out *in vivo* by administering the test compound and measuring its effect on respiration and/or body temperature.)

In addition, using cellular systems such as the two-hybrid system (Fields et al, Nature 340:245 (1989); Chien et al, Proc. Natl. Acad. Sci. USA 88:94=578 (1991)), proteins can be identified that interact with UCP2.

Using *in vivo* (or *in vitro*) systems, it may be possible to identify compounds that exert a tissue specific effect, for example, that increase UCP2 expression or activity only in fat or muscle or cells of the immune system.

Screening procedures such as those described above are useful for identifying agents for their potential use in pharmacological intervention strategies. Agents that enhance UCP2 expression or activity can be used to treat disorders such as, hyperinsulinemia, glucose intolerance, diabetes, obesity and syndrome X. Compounds that suppress UCP2 expression or inhibit its activity can be used to treat wasting associated, for example, with cancer, AIDS, cachexia and anorexia. Agents that suppress UCP2 expression or inhibit its activity can also be

used to induce hypothermia, for example, when  
advantageous in surgical settings, including  
transplantation. Such agents can also be used to  
block hyperthermia, for example, during thyroid storm.  
5 Compounds that enhance UCP2 expression or stimulate  
its activity can also be used to treat hypothermia.  
Given the similarity in the patterns of leptin and  
UCP2 expression, agents that modulate UCP2 can be  
expected to modulate leptin expression. However,  
10 leptin has been shown not to influence UCP2 expression  
in vitro or in vivo.

#### Pharmaceutical Compositions

The present invention also relates to  
15 pharmaceutical compositions comprising, as active  
agent, the proteins, peptides, nucleic acids or  
antibodies of the invention. The invention also  
relates to compositions comprising, as active agent,  
compounds selected using the above-described screening  
20 protocols. Such compositions include the active agent  
in combination with a pharmaceutically acceptable  
carrier. In the case of naked DNA, the "carrier" can  
be gold particles. In the case of compositions to be  
applied topically, the composition can take the form  
25 of a gel, cream, ointment or lotion. The amount of  
active agent in the composition can vary with the  
agent, the patient and the effect sought. Likewise,



the dosing regimen can vary depending on the composition and the disease/disorder to be treated.

Detection/Diagnosis:

5           The present invention further relates to methods of identifying individuals at increased risk for developing certain diseases/disorders, including hyperinsulinemia, glucose intolerance, type II diabetes, obesity, syndrome X, immunological  
10           dysfunction and body temperature dysfunction. One such method comprises: (a) obtaining from a mammal (e.g. a preobese human) a biological sample, (b) detecting the presence in the sample of a UCP2 gene product and (c) comparing the amount of the gene  
15           product present in the sample with that in a control sample. Advantageously, the biological sample is taken after the consumption of a high fat meal. In accordance with this method, the presence in the sample of altered (e.g. diminished) levels of UCP2  
20           gene product indicates that the subject is predisposed to the above-indicated diseases/disorders.

          Biological samples suitable for use in this method include biological fluids such as blood. Tissue samples (e.g. biopsies) can also be used in the  
25           method of the invention, including samples derived from muscle or fat. Cell cultures or cell extracts derived, for example, from tissue biopsies can also be used.

The detection step of the present method can be effected using standard protocols for protein/mRNA detection. Examples of suitable protocols include Northern blot analysis, immunoassays (e.g. RIA, Western blots, immunohistochemical analyses), and PCR.

The present invention also relates to methods of identifying individuals having elevated or reduced levels of UCP2, which individuals are likely to benefit from therapies designed to suppress or enhance UCP2 expression, respectively. As indicated above, a biological sample from a preobese subject can be screened for the presence of diminished levels of UCP2 gene product, particularly in response to high fat intake, the presence of depressed levels of the gene product, relative to a normal population (standard), being indicative of predisposition to obesity, type II diabetes or syndrome X. Such individuals would be candidates for anti-obesity therapy (e.g. treatment with appetite suppressants). The identification of elevated levels of UCP2 in a wasting patient (e.g. a cancer, AIDS or anorexia patient) would be indicative of an individual that would benefit from treatment with agents that suppress UCP2 expression or activity. The identification of low levels of UCP2 in a hypothermic patient or obese patient would be indicative of an individual that would benefit from agents that induce UCP2 expression or activity.

5 The present invention also relates to a kit that can be used in the detection of UCP2 expression products. The kit can comprise a compound that specifically binds UCP2 (e.g. binding proteins (e.g. antibodies or binding fragments thereof (e.g. F(ab')<sub>2</sub> fragments)) or UCP2 mRNA (e.g. a complementary probe or primer), for example, disposed within a container means. The kit can further comprise ancillary reagents, including buffers and the like.

10 The diagnostic methodologies described herein are applicable to both humans and non-human mammals.

#### Therapy:

15 The present invention relates to methods of treating diseases/disorders such as hyperinsulinemia, glucose intolerance, diabetes, obesity, syndrome X, cancer and hypothermia by increasing UCP2 activity and/or expression. The invention also relates to methods of treating inflammation, anorexia and wasting (cachexia) (e.g. associated with cancer or AIDS), of  
20 reducing fever and blocking hyperthermia (e.g. thyroid storm) and to methods of inducing hypothermia (eg when advantageous for surgery and transplant), by decreasing UCP2 activity and or expression. These  
25 methodologies can be effected using compounds selected using screening protocols such as those described above and/or by using the gene therapy and antisense approaches described below.

Gene therapy can be used to effect targeted expression of UCP2, for example, in fat tissue and muscle to reduce fat depots or in cancer cells to cause thermodestruction or metabolic collapse/death of the cells. The UCP2 coding sequence can be cloned into an appropriate expression vector and targeted to a particular cell type(s) to achieve efficient, high level expression. Introduction of the UCP2 coding sequence into target cells can be achieved, for example, using particle mediated DNA delivery (William et al, Proc. Natl. Acad. Sci. USA 88:2726 (1991); Haynes, Advances in Drug Delivery Reviews 21:3 (1996)), direct injection of naked DNA (Levy et al, Gene Therapy 3:21 (1996)), liposomal DNA transport (Felgner, Human Gene Therapy 1:1791 (1996)), or viral vector mediated transport (Smith et al, Gene Therapy 3:190 (1996)). Tissue specific effects can be achieved, for example, in the case of virus mediated transport by using viral vectors that are tissue specific, or by the use of promoters that are tissue specific (e.g. leptin and  $\alpha$ P2 promoters can be used to achieve expression in white adipose tissue and the myosin light chain kinase promoter can be used to achieve expression in skeletal muscle) (see also Warden et al, In Regulation of Body Weight: biological and behavioral mechanisms, C. Bouchard and G.A. Bray, eds. (West Sussex; John Wiley & Sons Ltd.), pp. 285-305). Combinatorial approaches can also be used to

ensure that the UCP2 coding sequence is activated in the target tissue (Butt et al, Gene Exp. 4:319 (1995); Hart, Semin. Oncol. 23:1521 (1996)).

5 Antisense oligonucleotides complementary to UCP2 mRNA can be used to selectively diminish or ablate the expression of the protein, for example, at sites of inflammation. More specifically, antisense constructs or antisense oligonucleotides can be used to inhibit the production of UCP2 in high expressing cells  
10 (spleen, thymus, leucocytes, bone marrow and stomach). Antisense mRNA can be produced by transfecting into target cells an expression vector with the UCP2 gene sequence, or portion thereof, oriented in an antisense direction relative to the direction of transcription.  
15 Appropriate vectors include viral vectors, including retroviral, adenoviral, and adeno-associated viral vectors, as well as nonviral vectors. Tissue specific promoters can be used (e.g. leptin gene promoter or aP2 gene promoter specific for adipose cells, muscle  
20 creatine kinase promoter specific for skeletal muscle and lymphoid cell promoters). Alternatively, antisense oligonucleotides can be introduced directly into target cells to achieve the same goal. (See also other delivery methodologies described above in  
25 connection with gene therapy.) Oligonucleotides can be selected/designed to achieve a high level of specificity. (See also Matteucci et al, Nature 384:20 (1996)).

5 It has been recently demonstrated that increased O<sub>2</sub> consumption associated with cachexia of malignancy can be attenuated by indomethacin, a cyclooxygenase inhibitor. This is thought to be due to inhibition of prostaglandin production (Roe et al, Metabolism 46:359 (1997)). Thus, agents that block UCP2 expression and/or activity can be expected to be useful in the treatment of cachexia.

10 The therapeutic methodologies described herein are applicable to both human and non-human mammals (including cats and dogs).

Certain aspects of the present invention are described in greater detail in the non-limiting Examples that follow.

#### 15 EXAMPLES

The following experimental methodologies relate particularly to Examples I and III-VII that follow.

20 Yeast transfections and functional analysis using flow cytometry. The diploid yeast (*Saccharomyces cerevisiae*) strain W303 (a/ $\alpha$ ; ade2-10; his 3-11-15; leu2-2, 112; ura3-1; can1-100; try- $\Delta$ 1) was used for expression of UCP2. The pYedP-UCP2 or UCPmut  
25 expression vectors were introduced into yeast by electroporation and transformants were selected for uracil auxotrophy as described by Bouillaud et al (EMBO J. 13:1990 (1994)). Expression of the UCP2,

UCP1 or UCPmut under the control of the gal-cyc promoter was induced by galactose in the absence of glucose. Growth of yeast for flow cytometry analysis and measurement of mitochondrial membrane potential using the DiOC6(3) fluorescent probe (3,3'-dihexyloxacarbocyanine iodine, Molecular Probes, Eugene, OR) are described by Bouillaud et al (EMBO J. 13:1990 (1994)).

**Isolation of UCP2 cDNA clones.** Human and mouse UCP2 IMAGE consortium clones (Washington University, St. Louis - Merck) identified in the EST database (<http://www.ncbi.nlm.nih.gov>) were purchased (Research Genetics, Huntsville, Alabama). Human UCP2 sequence was derived from IMAGE clones 129216, 158373, 163246 and 236034. GenBank accession for the human UCP2 coding sequence is U76367. Mouse UCP2 was also isolated independently (DR) by screening a mouse muscle cDNA library with rat UCP1 (GenBank accession U69135). More than 20 mouse and human UCP2 clones are currently in the EST database, while only one UCP1 clone (IMAGE 529189) is represented in the EST database.

**Mouse chromosomal mapping.** The chromosomal position was determined by linkage mapping of restriction fragment length polymorphisms (RFLPs) in an interspecific cross as previously described (Watson et al, Mamm. Genome 2:158 (1992)). Hybridization of human UCP2 probe (892 bp insert of IMAGE clone 129216)

defined EcoRI RFLPs that segregated in the strains used (C3H, 5.5 kb; *M. spretus*, 9.0 kb). Gene order was determined by analyzing all haplotypes and minimizing crossover frequencies. The reference loci in this interspecific cross that includes 2000 chromosomal markers are indicated with citations in an online database (<http://www.informatics.jax.org/crossdata.html>: select "seldin cross" and chromosome number 7).

#### EXAMPLE I

##### Sequence and Uncoupling Activity of UCP2

The amino acid sequence of human UCP2 is 59% identical to human UCP1 (UCP2 sequence shown in Figure 1A). The predicted coding sequence produces a protein of 309 amino acids with a molecular weight of 33,218 Daltons and an isoelectric point of 10.0. The amino acid sequence of mouse UCP2 (GenBank U69135) is 95% identical to human UCP2. Several protein motifs are conserved between UCP1 and UCP2. Both exhibit three mitochondrial carrier protein motifs, consistent with roles as ion transporters of the inner membrane, while the amino acids essential to ATP binding are also conserved.

To test the hypothesis that UCP2 is an uncoupling protein, yeast were transfected with UCP2 in an expression vector as previously reported for UCP1 (Bouillaud et al, EMBO J. 13:1990 (1994)). Rates of



growth in liquid medium of transformed yeast were measured in the presence of galactose, which induces expression. Instantaneous generation times were compared after induction of vector, UCP2, control (empty) and UCP1 expression vectors. The values for UCP2 were  $260 \pm 29$  minutes ( $\pm$ SD,  $n=5$ ), whereas the mean value for the strain transfected control vector (pYeDP vector) is  $188 \pm 13$  minutes ( $n=5$ ,  $p < 0.01$  versus UCP2) and the mean value for UCP1 is  $226 \pm 10$  minutes ( $n=10$ ,  $p < 0.01$  versus UCP2). These results indicate that UCP2 slows the growth of yeast more than does UCP1 (Bouillaud et al, EMBO J. 13:1990 (1994)).

The *in vivo* decrease of membrane potential associated with uncoupling of respiration was analyzed by flow cytometry of yeast labeled with the potential sensitive probe DiOC6(3). Using this procedure, it was previously demonstrated that UCP1 decreases the mitochondrial potential of transformed yeast (Bouillaud et al, EMBO J. 13:1990 (1994)).

Mitochondrial potential is strongly decreased in strains containing either the UCP2 or UCP1 expression vectors, in contrast to strains containing empty or UCPmut (A36P/K150L mutations that inhibit UCP1 activity) expression vectors (Figure 2). In the case of UCP2, 2 peaks were observed, one equivalent to cells expressing UCP1 and a second peak corresponding to a very low mitochondrial potential. Mean fluorescence intensities were: empty vector, 65.5;

SUBSTITUTE SHEET (RULE 26)

UCPmut, 43.0; UCP1, 22.5; UCP2, 15.5 and UCP2 + mCICCP, 0.4 (Figure 2). Very similar results were obtained in two additional experiments. Results of a representative experiment are shown in Figure 2. These data show that UCP2 influences mitochondrial activity and can partially uncouple respiration.

#### EXAMPLE II

##### Organization of Mouse and Human UCP2 Genes

*A map showing the organization of the mouse UCP2 gene present in MMU2-L2. The mouse DNA inserted in lambda phage is 13.9 kb long and contains all the 8 exons and introns, and 5.3 kb of DNA upstream of the putative transcriptional start site (+1 site). (See Fig. 7) The DNA has been sequenced from -934 to +8600 bp (Fig. 8B). The region -246/+1 of the mouse UCP2 gene was placed in front of a CAT reporter gene. This CAT construct exhibited a promoter activity when transfected in cultured adipocytes (see Example X). A fragment of 1.8 kb located in the 5' end of the gene (Fig. 8A) has also been sequenced.*

*A map showing the organization of the human UCP2 gene present in hUCP-g2. The human DNA inserted in lambda EMBL3 phage is 14 kb long (see Fig. 9). It contains all the 8 exons and introns, and a minimum of 3 kb of DNA upstream of the putative +1 site. Four regions, referred to as sequences 1, 2, 3 or 4 in*

Figures 10A-D., have been sequenced. Sequence 1 corresponds to 640 bp of DNA forming the 5' extremity of the human DNA. Sequence 2 corresponds to a 1161 bp DNA from positions bp -511 to +650. This fragment contains the putative proximal human UCP2 promoter.

Human and mouse UCP2 genes have a very similar organization, exons 1 and 2 being noncoding exons, whereas exons 3-6 are coding exons.

#### EXAMPLE III

##### Expression of UCP2 in Human and Mouse Tissues

To determine the possible sites of physiologic function of UCP2, Northern blot analyses were performed. UCP2 tissue distribution is markedly different from that of UCP1 (Figure 3A). Probing of a multiple tissue northern blot from pooled adult tissues reveals UCP2 mRNA of 1.6 kb size present in skeletal muscle, lung, heart, placenta and kidney. In mice, UCP2 is expressed in brown adipose tissue (BAT) as well as white adipose tissue (WAT), and at high levels in heart and kidney. Probing of a Human mRNA Master Blot (Clontech, Palo Alto, CA) with UCP2 reveals that it is expressed at high levels in skeletal muscle, spleen, thymus, leukocytes, bone marrow and stomach. Low levels of UCP2 mRNA expression were observed in liver or brain.

### EXAMPLE IV

## Mapping of UCP2 in Mice and Humans

Additional evidence for the role of UCP2 in obesity and diabetes was sought by determining the chromosomal position of UCP2. Whole genome mapping of mouse genetic crosses has identified at least 10 chromosomal loci influencing spontaneous and diet-induced obesity (Horvat et al, Genetics 139:1737 (1995); Collins et al, Mamm. Genome 4:454 (1993); Warden et al, J. Clin. Invest. 95:1545 (1995); West et al, J. Clin. Invest. 94:1410 (1994); Taylor et al Genomics 34:389 (1996); Seldon et al, J. Clin. Invest. 94:269 (1994)). Identification of the specific genes underlying these loci remains a challenge. A positional candidate approach, in which genes with suspected roles in obesity and insulin resistance are shown to have chromosomal loci co-incident with known obesity and diabetes loci, has been used to show that UCP2 is such a novel candidate gene. UCP2 (*Ucp2*) was mapped to mouse chromosome 7, tightly linked to the *tubby* mutation (Figure 3B). Two independent sequence tagged sites derived from human UCP2 clones (Expressed sequence tags, EST 143230 and 226515) have been mapped to the homologous region of the long arm of human chromosome 11, between D11S916 and D11S911 (WI-1672 and WI-13873, Whitehead Institute Center for Genome Research radiation hybrid panel, <http://www.ncbi.nlm.nih.gov/SCIENCE96/>), consistent

with mouse/human homology relationships suggesting an 11q13 chromosomal band location (DeBray et al, Genomics 33:337 (1996)).

5

**EXAMPLE V****UCP2 is a Positional Candidate Gene for Mouse and Human Obesity and Hyperinsulinemia Loci**

10

The chromosomal mapping of *UCP2* is co-incident with quantitative trait loci (QTLs) for obesity from at least three independent mouse models, one congenic strain, and human insulin dependent diabetes locus-4 (IDDM4) (Warden et al, J. Clin. Invest. 95:1545 (1995); Taylor et al, Genomics 34:389 (1996); Seldin et al, J. Clin. Invest. 94:269 (1994); Hashimoto et al, Nature 371:61 (1994)). Diet-induced obesity and diabetes have been demonstrated in the C57BL/6J (B6) mouse, while the A/J strain is resistant to the high-fat diet (Surwit et al, Diabetes 37:1163 (1988)). This syndrome is not due to inactivity (Brownlow et al, Physiology & Behavior 60:37 (1996)) or to hyperphagia (Surwit et al, Metabolism 44:545 (1995)) as B6 mice show higher feed efficiency than A/J mice (weight gained/calories consumed), when fed a high fat diet. Whole genome mapping has shown that a QTL for plasma leptin, insulin, and glucose is located on mouse chromosome 7, peaking near the *tubby* locus (Surwit et al, Diabetes 45:78A (1996)). Sequence analysis and mRNA expression levels suggest that *tubby*

15

20

25

is not the underlying gene for this QTL. When Northern blots of white adipose tissue from A/J and B6 mice were hybridized with the UCP2 probe, mRNA levels were found to be higher in the obesity-resistant A/J than the obesity prone B6 (Figure 4). In addition, UCP2 mRNA levels were significantly increased by the high-fat diet in A/J mice at early time points in the diet (days 7, 18 and 25), but only modestly in B6 animals (Figure 3C). Core body temperature measured by a rectal temperature probe is also increased in A/J and B6 mice consuming the high fat diet, but to a greater extent (2X) in A/J than in B6 mice. These data are highly significant (n=10 for each group,  $p < 0.01$ ). However, after progression to severe obesity in the B6 mouse on a high fat diet for 18 weeks, levels of UCP2 were dramatically increased in both strains, but in contrast to early time points (7, 18, 25 days) levels of UCP2 were greater in the diet-induced obese B6 mice relative to the leaner A/J strain. Additionally, core body temperature increased more in A/J mice than in B/6 mice following the introduction of a high fat diet ( $p < 0.05$ ). Similar over expression of UCP2 is observed in adult, genetically obese ob/ob and db/db mice.

Regulation of UCP2 has been examined for factors that would regulate UCP1. In 11-week-old fed male Swiss mice, UCP2 mRNA was detected in BAT, WAT, thigh muscle and liver. Exposure at 4°C or treatment with

the  $\beta 3$  adrenergic agonist CL316,243 for 10 days did not influence UCP2 mRNA levels in these tissues. Finally, administration of leptin (2  $\mu$ g, 2x/day for 4 days) failed to increase UCP2 expression. Leptin has been shown to increase sympathetic outflow to brown fat, thus potentially increasing UCP2 expression. Thus, UCP2 is regulated very differently than UCP1, which is induced by cold and  $\beta 3$  agonists. However, these data do not rule out the possibility that the sympathetic nervous system regulates UCP2.

#### EXAMPLE VI

##### **Increase in UCP2 Expression by Thiazolidinediones**

Thiazolidinediones are known insulin-sensitizing agents that lower plasma glucose levels, and long chain fatty acids have been shown to be ligands for the PPAR (peroxisome proliferator activated receptor) family of receptors. Accordingly, a study was undertaken to determine whether agents that stimulate PPAR $\gamma$  could increase expression of UCP2 in a model adipocyte cell line, HIB-1B.

HIB 1B cells were grown for 7 days in DMEM + 10% charcoal-stripped serum (Media) or with the addition of the thiazolidine dione BRL 49653 (1 $\mu$ M) and the RXR $\alpha$  ligand LGD-1069 (0.1 $\mu$ M) (TZD/RXR). The results are the average of 2 independent experiments of 4 samples each. \*, significantly different from Media samples,  $p < 0.001$ . (See Fig. 6).

EXAMPLE VII**Respiratory Rates**

Respiratory rates have been measured in mitochondria isolated from yeast containing the empty pYeDP vector and the pUCP2 vector (Table 1). Upon addition of the FCCP uncoupler, the respiratory rate was increased by a factor of 7.1 in control mitochondria, while it was only increased by a factor of 4.3 in pUCP2 mitochondria (Table 1). This result clearly demonstrates that UCP2 has uncoupling activity. Addition of ADP stimulates respiration more in control mitochondria, again showing the UCP2 mitochondria are more uncoupled.

Table 1

	Basal rate	Uncoupled		ADP addition	
		rate	fold	rate	fold
pYeDP	14	100	7.1	33	2.3
pUCP2	23	100	4.3	42	1.8

EXAMPLE VIII**Role of UCP2 in a mouse model of spontaneous obesity**

The *Ucp2* gene is co-incident with a QTL for spontaneous multifactorial obesity in BSB mice. BSB mice are derived from a backcross of (C57BL/6J x *Mus spretus*) F1s x C57BL/6J. A locus with significant QTLs for body fat percent, hepatic lipase activity and



plasma cholesterol is located on distal mouse chromosome 7 surrounding the *tubby* locus. mRNA extracted from the spleens of 16 BSB mice was probed on northern blots for UCP2 and actin. Levels of mRNA were quantitated by phosphorimager and UCP2 was normalized with actin. A negative correlation of UCP2/actin mRNA levels with body fat ( $p=0.016$ ) and HDL cholesterol ( $p=0.023$ ) was observed. The hypothesis that body temperature is correlated with body fat in BSB mice was also investigated. Core body temperatures were determined by rectal thermometers every week after weaning of 45 BSB mice. These BSB mice demonstrated a negative correlation of body temperature with percent body fat at sacrifice. P-values for the correlation of were  $p=0.018$  at 6 weeks old ( $r^2 = 0.12$ ) and  $p=0.09$  at week 8 ( $r^2=0.10$ ). These results indicate that UCP2 influences body fat percent by altering body temperature in a model of spontaneous obesity exhibited by mice maintained on a chow diet. (See Fisler et al, Obesity Res. 1:271 (1993); Warden et al, J. Clin. Invest. 92:773 (1993); Warden et al, J. Clin. Invest. 95:1545 (1995).)

#### EXAMPLE IX

##### Common Sequence Polymorphisms of Human UCP2

Human mRNA was prepared from buffy coats (white blood cells) from 8 Hispanics who either have Type II diabetes or are the children of Type II diabetics.

The complete human UCP2 coding sequence was amplified by RT-PCR of mRNA using primer pair 3 (hUCP2.cds3) as described in the Brief Description of Fig. 1E. The products were cloned into the Invitrogen pCR2.1 vector as described by the manufacturer. Sequencing was performed with 8 primers on each of the clones. Sequences were compared with the hUCP2 sequence submitted to Genbank (Accession No. U76367). 5 polymorphisms were observed in at least 2 of the 8 people sequenced. Each of these confirmed polymorphisms was also observed with all overlapping primers. 5 additional polymorphisms were observed with only one person or have conflicting data from different primer pairs. (See Table 2) Restriction enzymes have been identified that would differentially digest each of the alleles.

TABLE 2

Polymorphism	1	2	3	4	5
base number in terms of entire dna sequence	445	502	508	869	913
amino acid number in terms of open reading frame seq.	34	53	55	104	190
hUCP2.genbank	L	A	A	K	N
Confirmed polymorphisms, observed at least twice.	L	A	V	K	S
Polymorphisms observed only once.	P	V	V	K	S
Short name			A55V		N190S

Polymorphism	6	7	8	9	10
base number in terms of entire dna sequence	1052	1068	1142	1224	1262
amino acid number in terms of open reading frame seq.	236	242	266	294	306
hUCP2.genbank	D	Y	P	L	E
Confirmed polymorphisms, observed at least twice.	D	Y	P	M	E
Polymorphisms observed only once.	E	N	P	M	E
short name				L294M	

*Restriction enzymes of HUCP2.genbank for polymorphism analysis:*

Polymorphism 1: Frequency of base change T to C at 445 of entire expressed is 77% T and 33% C. Amino acid change occurs in polymorphisms observed only once from L to P.

site= t/ccgga, possible enzymes= AccIII, BseAI, BseAI, BsiMI, Bsp13I, BspEI, Kpn2I, MroI, There are no other areas where this enzyme would cut.

Polymorphism 2: Frequency of base change C to T at 502 of entire expressed sequence is 22% C, 88% T; base change T to A at 503 of entire expressed sequence is 64% T, 36% A. Amino acid change occurs in edited all sequence from A to V. Restriction enzyme cleavage site= /gtnac, possible enzyme is MaeIII. Other areas that this enzyme cuts are: 119, 583, 865.

Polymorphism 3: Frequency of base change C to T at 508 of entire expressed sequence is 57% C and 43% T as shown in 2 primers. Amino acid change occurs in edited and edited all sequences from A to V. Restriction enzyme cleavage site=gcn/ngc, possible enzyme is Cac8I. Other areas that this enzyme cuts are: 58, 70, 153, 250, 254, 336, 340, 364, 670, 756, 901.

Polymorphism 4: Frequency of base change C to T at 869 of entire expressed sequence is 73% C and 27% T, shown in 2 primers. Seventy-five percent of examined individuals are T. No amino acid change occurs at this possible polymorphism site. Restriction enzyme cleavage site= cctc, possible enzyme is MnlI. Other areas that this enzyme cuts are: 97, 231, 278, 357, 546, 594, 609, 624, 639, 695, 796, 824, 891.

Polymorphism 5: Frequency of base change A to G at 913 of entire expressed sequence is 80% A and 20% G as

shown in two primers. Twenty-five percent of examined individuals are G. Amino acid change occurs in edited and edited all sequences from N to S. Restriction enzyme cleavage site=cmg/ckg where m=a,c and k=g,t; possible enzymes= MspAII, NspBII or site= cag/ctg with enzyme PvuII. Other areas that cmg/ckg cuts are:cat/ctg: 65, 877; ccg/cgg: 177.

Polymorphism 6: Base change C to A at 1052 of entire expressed sequence is included in edited all sequence because 4 out of 8 exhibit A in primer -21M13 and C in primer 241.1. Amino acid change occurs in edited all sequence from D to E. Restriction enzyme cleavage site: gacn/nngtc, possible enzymes are AspI, AcsI, Tth111I. There are no other areas where this site occurs.

Polymorphism 7: Base change T to A at 1068 of entire expressed sequence is included in edited all sequence because 5 out of 8 exhibit A in primer -21M13 and T in primer 241.1. Four out of five individuals are the same ones changing as in Polymorphism 6. Amino acid change occurs in edited all sequence from Y to N. There are no enzymes to cleave near this possible polymorphism.

Polymorphism 8: Frequency of base change C to T at 1142 of entire expressed sequence is 64% C and 36% T

as shown in 4 primers. No amino acid change occurs at this possible polymorphism site. Restriction enzyme cleavage site= t/cga, possible enzymes are TagI and TthHB8I. There are no other areas where this site occurs.

Polymorphism 9: Frequency of base change C to A at 1224 of entire expressed sequence is 64% C and 36% A as shown in 4 primers. The same individuals have the change from the expressed sequence as in Polymorphism 8. An amino acid change occurs in edited and edited all sequences from L to M. Restriction enzyme cleavage site=gcagc, possible enzymes are BbI and Bst71I. Other areas that this enzyme cuts are 258 and 368.

Polymorphism 10: Frequency of base change G to A at 1262 of entire expressed sequence is 57% G and 43% A. Six out of seven exhibit A. No amino acid change occurs at this possible polymorphism site. Restriction enzyme cleavage site= cctc, possible enzyme is MnlI. The enzyme cleaves the complementary DNA but not the hUCP2.GenBank sequence at this site. Other areas that this enzyme cuts are 97, 231, 278, 357, 522, 536, 594, 609, 624, 639, 695, 796, 824, 891.

*Amplification of human UCP2 exons containing polymorphisms.*

The following primers are designed to amplify hUCP2 exons 4, 6, 7 and 8 from genomic DNA. Common amino acid variants are present in exons 4, 6, and 8: A55V is in exon 4, N190S is in exon 6, and L294M is in exon 8. N190S is expected to alter a PvuII site. It is expected that Ser 190 will cut and Asn190 will not cut with PvuII. Primers for exon 7 have also been designed.

Results:

Exon 4: Both primer pairs 1 and 2 produce single bands of the expected size.

Exon 6: Primer pair 2 (3219 - 3399) produces a single band of the expected size. Primer pair 1 (3147- 3416) did not work.

Exon 7: Primer pair 1 (4316-4594) produces a single band of the expected size. Primer pair 2 (4284 - 4598) did not work.

Exon 8: Primer pairs 1 and 2 both produce multiple bands, including ones of the expected size.

PCR conditions: Clontech KlenTaq plus polymerase is used. PCR conditions were 94°C 1 min, 94°C 30 sec, 66°C 30 sec, 68°C 30 sec, repeat steps 2-4 29-times, 4°C indefinite. Reaction volume was 25 µl in an MJ research PTC-200.

# hUCP2g.e4f1

5

```
>> Melting Temperature:      60.5
```

```
>> GC Clamp Strength: 57
```

10

```
>> Primer Dimer:      4
```

hUCP2g.e4r1

15

```
>> Melting Temperature:      60.6
```

```
>> GC Clamp Strength: 45
```

20

```
>> Primer Dimer:      6
```

```
>> Product Size: 362
```

25

11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 10

hUCP2g.e4f2

30

```
>> Melting Temperature:      59.6
```

>> GC Clamp Strength: 47

35

>> Primer Dimer: 4

hUCP2g.e4r2

```
>> Length: 18
```

```
>> Max AT Run Length: 3
```

>> GC Clamp Strength: 60



```

>> % GC Content:    61
>> Hairpin Stem:    2
>> Primer Dimer:    6
>> No Palindromes.
5  >> Product Size:    424
>> Annealing Temperature:    59.9    (Try T = 54.4)
=====
amplifies bp 3147-3416 of hUCP2g, Meudon
hUCP2g.e6f1
10 Forward Oligo:    5'>    CAAGACCATT GCCCGAAG    <3'
>> Length:    18
>> Melting Temperature:    60.2
>> Max AT Run Length:    3
>> GC Clamp Strength:    52
15 >> % GC Content:    56
>> Hairpin Stem:    2
>> Primer Dimer:    6
>> No Palindromes.
hUCP2g.e6r1
20 Reverse Oligo:    5'>    ATGGGGGAAG GGTGAGAC    <3'

>> Length:    18
>> Melting Temperature:    59.7
>> Max AT Run Length:    2
25 >> GC Clamp Strength:    14
>> % GC Content:    61
>> Hairpin Stem:    2
>> Primer Dimer:    4
>> No Palindromes.
30 >> Product Size:    269
>> Annealing Temperature:    59.2    (Try T = 54.7)
=====
amplifies bp 3219-3399 of hUCP2g, Meudon
hUCP2g.e6f2
35 Forward Oligo:    5'>    TTCCTCCTCC CCGATACTC    <3'
>> Length:    19
>> Melting Temperature:    59.0
>> Max AT Run Length:    3
>> GC Clamp Strength:    8
>> % GC Content:    58
>> Hairpin Stem:    2
>> Primer Dimer:    4

```

```

>> No Palindromes.
hUCP2g.e6r2
Reverse Oligo: 5'> CCCAGCACCG TCTACCTC <3'
>> Length: 18
5 >> Melting Temperature: 59.2
>> Max AT Run Length: 2
>> GC Clamp Strength: 21
>> % GC Content: 67
>> Hairpin Stem: 2
10 >> Primer Dimer: 4
>> No Palindromes.
>> Product Size: 180
>> Annealing Temperature: 58.1 (Try T = 54.0)
=====
15 amplifies bp 4840-5338 of hUCP2g, Meudon
hUCP2g.e8f1
Forward Oligo: 5'> AAGTGGGAGG TGGAGGTG <3'
>> Length: 18
>> Melting Temperature: 58.4
20 >> Max AT Run Length: 2
>> GC Clamp Strength: 29
>> % GC Content: 61
>> Hairpin Stem: 0
>> Primer Dimer: 4
25 >> No Palindromes.
hUCP2g.e8r1
Reverse Oligo: 5'> AGCTACAAGA GAGGAGGAGA CG <3'
>> Length: 22
>> Melting Temperature: 59.3
30 >> Max AT Run Length: 2
>> GC Clamp Strength: 30
>> % GC Content: 55
>> Hairpin Stem: 1
>> Primer Dimer: 4
35 >> Longest palindrome has 4 bases (bases 1 to 4).
>> Product Size: 520
>> Annealing Temperature: 59.5 (Try T = 53.4)
=====
amplifies bp 4831-5270 of hUCP2g, Meudon
hUCP2g.e8f2
Forward Oligo: 5'> CAGTGAGGGA AGTGGGAGG <3'
>> Length: 19

```

49

```

>> Melting Temperature:    60.7
>> Max AT Run Length:      2
>> GC Clamp Strength:      57
>> % GC Content:           63
5  >> Hairpin Stem:          1
>> Primer Dimer:           2
>> No Palindromes.
hUCP2g.e8r2
10 Reverse Oligo:    5'>    GGGGCAGGAC GAAGATTC    <3'
>> Length:          18
>> Melting Temperature:    60.6
>> Max AT Run Length:      3
>> GC Clamp Strength:      19
15 >> % GC Content:          61
>> Hairpin Stem:          2
>> Primer Dimer:          6
>> No Palindromes.
>> Product Size:          457
>> Annealing Temperature:    60.1    (Try T = 55.6)
20 =====
amplifies 4316-4594 of hUCP2g, Meudon
hUCP2g.e7f1
Forward Oligo:    5'>    CTTGCCTGCT CCTCCTTG    <3'
25 >> Length:              18
>> Melting Temperature:    59.7
>> Max AT Run Length:      2
>> GC Clamp Strength:      35
>> % GC Content:           61
30 >> Hairpin Stem:          1
>> Primer Dimer:          6
>> No Palindromes.
hUCP2g.e7r1
Reverse Oligo:    5'>    GGTGGTTCTC TCCCACCC    <3'
35 >> Length:              18
>> Melting Temperature:    60.3
>> Max AT Run Length:      2
>> GC Clamp Strength:      57
>> % GC Content:           67
>> Hairpin Stem:          3
>> Primer Dimer:          6
>> No Palindromes.
>> Product Size:          278

```

SUBSTITUTE SHEET (RULE 26)

50

```

>> Annealing Temperature:    60.4    (Try T = 54.7)
=====
amplifies 4284-4598 of hUCP2g, Meudon
hUCP2g.e7f2
5 Forward Oligo: 5'> CCTTCTGGAA TGATGGGTG <3'
>> Length: 19
>> Melting Temperature: 58.9
>> Max AT Run Length: 3
>> GC Clamp Strength: 45
10 >> % GC Content: 53
>> Hairpin Stem: 2
>> Primer Dimer: 6
>> No Palindromes.
hUCP2g.e7r2
15 Reverse Oligo: 5'> GAACTGGGTG GGGAGGAC <3'
>> Length: 18
>> Melting Temperature: 60.3
>> Max AT Run Length: 2
>> GC Clamp Strength: 26
20 >> % GC Content: 67
>> Hairpin Stem: 2
>> Primer Dimer: 6
>> No Palindromes.
>> Product Size: 265
25 >> Annealing Temperature: 60.3 (Try T = 53.9)
=====

```

EXAMPLE X

## Immune Cell Regulation of UCP2

30 (A) A tumor mast cell line, the Rat Basophilic Leukemia cell (RBL), is a model system for studies of immune system signal transduction. In the RBL cell, binding of an antigen to the IgE receptor on the cell surface results in the activation of phospholipase C-gamma (PLCg). This event leads to the activation of protein kinase C (PKC) and an increase in intracellular calcium which are required for

SUBSTITUTE SHEET (RULE 26)

degranulation and gene expression in RBL cells. Several drugs can mimic the intracellular events and degranulation occurring in this cell type in response to antigen activation of the IgE receptor. The inophore Ionomycin causes the release of calcium from intracellular calcium stores and the phorbol ester PMA is able to directly activate PKC. Both of these drugs are required to achieve maximal activation of RBL cells. In order to determine the expression level of UCP2 in RBL cells, these cells were treated with Ionomycin (5mM), PMA (5mM) or both drugs. A Northern blot of the RNA from these cells revealed a ~4 fold induction of UCP2 expression in treated cells compared to untreated control cells. Addition of Ionomycin, PMA or both drugs simultaneously resulted in the same ~4 fold increase in UCP2 expression. Since PMA and Ionomycin treatment do not cause an additive or synergistic expression of UCP2, maximal activation of RBL cells may not be required for UCP2 gene expression.

(B) Bacterial lipopolysaccharide (LPS) regulates UCP2 mRNA in macrophages. Bacterial LPS is known to induce many of the features of infection, including fever. Thioglycolate elicited mouse macrophage (peritoneal exudate cells, PEC) have been studied for regulation of UCP2 mRNA by LPS. It has been shown that PEC express UCP2 and that LPS decreases expression by several fold over a 24 hour time course.

A study has also be made of UCP2 expression in cultured mouse macrophage RAW cells. These cells express UCP2. Addition of 100 ng/ml LPS in 5% calf serum decreases UCP2 mRNA levels by several fold after 24 hours. Lower doses of LPS are less effective, while shorter times produce smaller decreases. LPS seems to be more effective at decreasing UCP2 in cells maintained in 0.5% serum than in 5% serum. The LPS mediated decrease of UCP2 is not affected by indomethacin or by PGE2. All fo the macrophage northernns are normalized for actin expression and have been repeated at least once. These data show that LPS, which is known to influence immune function and to promote fever, can also influence immune function and to promote fever, can also influence UCP2 mRNA levels. The results are consistent with a role of UCP2 in immune function.

#### EXAMPLE XI

##### UCP2 Promoter Activity

A 246 bp putative promoter region for mouse UCP2 was cloned upstream of the chloramphenicol acetyl transferase gene in either the "sense" (also called "+" or "forward" orientation) or the "antisense" (or "-" or "reverse" orientation). When transfected into differentiated mouse HIB-1B adipocytes CAT activity (conversion of substrate into product) is observed for

the "sense" ("+") orientation, but not for the  
"antisense" ("-") orientation. (See Fig. 11).

#### EXAMPLE XII

##### The Linkage Study

###### Experimental Details:

Sample. A total of 640 individuals (299 males and 341 females) from 155 pedigrees were available for the present study. These were randomly ascertained families of French descent living in the Québec city area and recruited to participate in the Québec Family Study, a population-based study of the genetics of physiological fitness and body composition. The age of individuals in the sample ranges from 18 to 94 years.

Phenotypes. BMI was obtained from height and weight measurements (BMI=weight in kg/height in m). FM, fat-free mass and %FAT were determined from body density measurements obtained by underwater weighing using the conversion factor of Siri (Adv. Biol. Med. Phys. 4:239 (1976)). RMR was determined by indirect calorimetry measurements using an open-circuit indirect calorimeter with the ventilated hood technique (Derioz et al, J. Clin. Invest. 93:838 (1994)). Measurements were taken in the morning after an overnight fast, while subjects sat quietly in a semireclined position

for the 30 minute measurement period. The last 10 minutes were kept for calculation of the RMR. The  $VO_2$  and  $VCO_2$  data were converted into energy as recommended by Weir (J. Physiol. (Lond) 109:1 (1949)). The phenotypes were adjusted by sex, for age and age by regression procedures and RMR was further adjusted for FM and FFM. The residuals from the regressions were used for linkage analysis.

*DNA typing.* Genomic DNA was prepared from permanent lymphoblastoid cells by the proteinase K and phenol/chloroform technique. DNA was dialysed four times against TE buffer (10 mM Tris, 1 mM EDTA pH 8.0) for 6 hours at 4°C and ethanol precipitated.

Amplification (EasyCycler, Ericomp, San Diego, CA) was done in 96 wells microtiter plaques using 250 ng of genomic DNA 0.1 pmoles (D11S1321, D11S916) or 0.25 pmoles (D11S911) of the forward primer coupled to the infrared tag IRD41 (Licor) and, respectively, 0.1 or 0.4 pmoles of the reverse primer, 125  $\mu$ M dNTP's, and 0.3 U Taq polymerase (Perkin Elmer, Roche Molecular Systems, Branchburg, NJ) in PCR buffer (100 mM Tris-HCl pH 8.3, 15 mM  $MgCl_2$ , 0.5 M KCl, 0.01% gelatin) for a final volume of 10  $\mu$ l. PCR cycles consisted of 1 cycle at 93°C for 5 min., 10 cycles at 94°C for 20 sec, 57°C for 60 sec, and 24 cycles at 94°C for 20 sec., 52°C for 60 sec., except for D11S911 for which the first annealing temperature was set at 55°C, PCR



products were analyzed on automatic DNA sequencer (Li-Cor) using 18 cm glass plates. Typing was done assisted by computer (One Dscan, Scannalytics).

5        *Linkage analysis.* Relative-pair based methods were used to test for linkage between the phenotypes and the marker loci. In the presence of linkage between a marker locus and a quantitative trait (Y), relative pairs sharing a greater proportion of alleles  
10       identical by descent ( $\pi$ ) as the marker locus tend to have more similar phenotypes than pairs who share fewer alleles. Thus, under the hypothesis of linkage, a negative relationship is expected between  $\pi$  and the within pair variance. The sib-pair linkage method  
15       described by Haseman and Elston is the most widely used method to investigate linkage between a quantitative phenotype and a marker locus. This method has been extended to other types of relative pairs. Tests of linkage which combine the information  
20       from different types of relative pairs were developed and these tests have been shown to be more powerful than the Haseman and Elston method based only on sib-pairs. This relative pair linkage analysis has been implemented in the program RELPAL which considers  
25       the following five types of relative pairs: sibling, half-sibling, grandparent-grandchild, avuncular and first degree cousins. For each relative pair type, the statistic for testing linkage is obtained by

dividing the estimated regression coefficient ( $\beta$ ) by its standard error. Because the number ( $n$ ) of relative pairs could vary among the different types of relatives depending on the complexity of the pedigrees, the contribution of each type of relative pairs needs to be weighted in the overall linkage statistic which combine information from all relative pairs. The linkage test implemented in RELPAL is:

$$T = \frac{c^T \beta}{\sqrt{c^T \text{Var}(\beta) c}}$$

where  $\beta$  is a vector containing the  $\beta$ 's for each of the 5 types of relatives and  $c^T$  is a weighing vector based on  $n$  and the variance of  $\pi$  and equal to:  $(\text{Var}(\pi_s)n_s, \text{Var}(\pi_h)n_h, \text{Var}(\pi_g)n_g, \text{Var}(\pi_a)n_a, \text{Var}(\pi_c)n_c)$ , where subscripts  $s, h, g, a$  and  $c$  stand for siblings, half-siblings, grandparent-grandchild, avuncular and first degree cousins, respectively. In the Québec Family Study, since there are no half-sibs, only sibling, avuncular, grandparental and cousin pairs were used in the relative-pair linkage analysis.

#### Results:

A study was undertaken to investigate the potential linkage relationships between three microsatellite markers which encompass the UCP2 gene location on 11q13 with resting metabolic rate (RMR), body mass index (BMI), percentage body fat (%FAT) and fat mass (FM) in 640 individuals from 155 randomly

ascertained pedigrees from the Québec Family Study. Using a linkage analysis strategy based on sibling, avuncular, grandparental and cousin pairs, strong evidence of linkage was found between the marker D11S911 ( $p=0.000002$ ) and RMR with more moderate evidence for D11S916 ( $p=0.006$ ) and D11S1321 ( $p=0.02$ ). Suggestive evidence of linkage was also observed between D11S1321 and %FAT ( $p=0.04$ ) and FM ( $p=0.02$ ). It is concluded that the three markers encompassing the UCP2 locus and spanning a 5 cM region on 11q13 are linked to resting energy expenditure in adult humans.

The mouse *ucp2* gene was recently mapped to chromosome 7, closely linked to the *tubby* mutation, a mutation known to be responsible for the adult-onset obesity in this mouse model. Furthermore, the UCP2 mRNA level was found to be higher in the A/J mouse strain, which is resistant to diet-induced obesity, than in the obesity prone C5/BL/6J mouse. The evidence accumulated thus far on animal models indicates that the UCP2 gene plays a role in the development of obesity because of its role in energy metabolism. The human UCP2 gene has been mapped to chromosome 11q13 at a location distinct from *tubby* (11p15.1) but in the same chromosomal location as the Bardet-Biedl Syndrome locus (Online Mendelian Inheritance in Man, OMTM, Center for Medical Genetics, Johns Hopkins Univ., (Baltimore, MD) and National Center for Biotechnology Information, National Library

of Medicine (Bethesda, MD) (1986), WWW URL:  
<http://www3.ncbi.nlm.nih.gov/omim/>), a Mendelian  
Syndrome exhibiting obesity as one of its clinical  
features. UCP2 is also in the proximity (about 15 cM)  
of a locus (11q21-q22) recently uncovered through a  
genome-wide search and found to be linked to percent  
body fat in Pima Indians. To test the hypothesis that  
markers around the UCP2 gene may exhibit a linkage  
relationship with metabolic rate and body fat  
phenotypes, three markers (D11S916, D11S1321 and  
D11S911) were typed on 640 individuals from 155  
pedigrees from the Québec Family Study. Linkage  
studies were undertaken with RMR, BMI, %FAT and FM  
using four types of relatives. RMR was adjusted for  
the effects of age, sex, FM and fat-free mass, whereas  
BMI, %FAT and FM were adjusted only for age and sex  
effects.

Table 3 presents the linkage results with the  
number of relative pairs available in each case.  
Strong evidence of linkage was observed between the  
D11S911 and RMR while more moderate evidence for  
linkage was found for the other two markers. As there  
were few pairs of first degree cousins, the same  
analysis was repeated after removing these pairs. In  
the latter case, the p-value for linkage between  
D11S911 and RMR (n=212 pairs) was reduced to p=0.001,  
while the one between D11S916 and RMR (n=279 pairs)  
went from p=0.006 to p=0.02. Evidence of linkage

between D11S1321 and %FAT ( $p=0.04$ ) and FM ( $p=0.02$ ) was also found. No linkage was observed between the three markers and fat-free mass in this population.

Overall, these results indicate that the UCP2 gene which is encoded within the 5 cM span covered by these markers, plays a role in determining resting energy expenditure in humans.

Table 3. Relative-pair linkage analysis of resting metabolic rate and body fat variables with markers encompassing the UCP2 gene in the Québec Family Study

MARKER	HZ	cM	RMR	BMI	%FAT	FM
D11S916	0.72	78	0.006 (301)	0.27 (415)	0.50 (304)	0.11 (304)
D11S1321	0.64	79	0.02 (380)	0.36 (537)	0.04 (383)	0.02 (383)
D11S911	0.85	83	0.000002 (240)	0.38 (324)	0.23 (243)	0.26 (243)

Based on four different types of relative pairs: siblings (165-275 pairs); avuncular (47-134 pairs); grandparental (45-94 pairs); first degree cousins (22-34 pairs). The entries are the p-values with total number of relative pairs given in parentheses. HZ: heterozygosity; cM; distance in centimorgan.

\* \* \*

All documents cited herein are incorporated in their entirety by references.

What is claimed is:

1. A method of treating a disease or disorder associated with diminished UCP2 expression or activity comprising administering to an individual in need of such treatment an amount of an agent that enhances UCP2 expression or activity sufficient to effect said treatment.
2. The method according to claim 1 wherein said disease or disorder is obesity, diabetes, syndrome X, or hypothermia.
3. The method according to claim 1 wherein said disease or disorder is hyperinsulinemia or glucose intolerance.
4. The method according to claim 1 wherein said agent is an expression construct comprising a UCP2 encoding sequence or portion thereof that encodes functional UCP2.
5. A method of treating a disease or disorder associated with elevated UCP2 expression or activity comprising administering to an individual in need of such treatment an amount of an agent that inhibits UCP2 expression or activity sufficient to effect said treatment.

6. The method according to claim 5 wherein said disease or disorder is wasting, anorexia, inflammation, cachexia, fever, or hyperthermia.
7. The method according to claim 5 wherein said agent is an oligonucleotide that specifically hybridizes to UCP2 mRNA or an antibody, or binding fragment thereof, that is specific for UCP2.
8. A method of screening a compound for its ability to modulate UCP2 expression or activity comprising contacting a cell that expresses UCP2 with said compound under conditions such that modulation of UCP2 expression or activity can occur and measuring UCP2 expression or activity in the presence and absence of said compound and determining whether modulation is effected.
9. Isolated mammalian UCP2, or fragment thereof of at least 5 amino acids.
10. The UCP2 according to claim 9 wherein said UCP2 is human UCP2.
11. The UCP2 according to claim 10 wherein said UCP2 has the sequence shown in Figure 1A.



13. The nucleic acid according to claim 12 wherein said nucleic acid encodes human UCP2.

14. The nucleic acid according to claim 13 wherein said human UCP2 has the sequence shown in Figure 1A.

15. A recombinant molecule comprising a vector and the nucleic acid according to claim 12.

16. A host cell comprising the recombinant molecule according to claim 15.

17. A method of making UCP2 comprising incubating the host cell according to claim 16 under conditions such that said nucleic acid is expressed and said UCP2 is thereby produced.

18. An antibody specific for UCP2 or antigen binding fragment thereof.

19. An antisense construct comprising a vector and a nucleic acid operably linked to a promoter, wherein said nucleic acid is oriented with respect to the

promoter such that the transcript of the nucleic acid selectively hybridizes to UCP2 mRNA.

20. A host cell comprising the construct according to claim 19.

21. The cell according to claim 20 wherein said cell is a mammalian cell.

Fig. 1A.

DEFINITION hUCP2, 309 amino acids.  
 ORIGIN

```

1  MVGFKATDVP PTATVKFLGA GTAACIADLI TFPLDTAKVR LQIQGESQGP
51 VRATASAQYR GVMGTILTMV RTEGPRSLYN GLVAGLQRM SFASVRIGLY
101 DSVKQFYTKG SEHASIGSRL LAGSTTGALA VAVAQPTDVV KVRFQAQARA
151 GGGRRYQSTV NAYKTIAREE GFRGLWKGTS PNVARNAIVN CAELVTYDLI
201 KDALLKANLM TDDLPCHEFIS AFGAGFCTTV IASPDVVKT RYMNSALGQY
251 SSAGHCALTM LQKEGPRAFY KGFMPSFLRL GSNVVVMFVT YEQLKRALMA
301 ACTSREAPF
  
```

Fig 1B.

DEFINITION hUCP2 coding sequence, 930 bases including the stop  
 codon.  
 ORIGIN

```

1  ATGGTTGGGT TCAAGGCCAC AGATGTGCCC CTTACTGCCA CTGTGAAGTT
51 TCTTGGGGCT GGCACAGCTG CCTGCATCGC AGATCTCATC ACCTTTCCTC
101 TGGATACTGC TAAAGTCCGG TTACAGATCC AAGGAGAAAG TCAGGGGCCA
151 GTGCGCGCTA CAGCCAGCGC CCAGTACCGC GGTGTGATGG GCACCATCTT
201 GACCATGGTG CGTACTGAGG GCGCCCGAAG CCTCTACAAT GGGCTGGTTG
251 CCGGCCTGCA GCGCCAAATG AGCTTTGCCCT CTGTCCGCAT CGGCCCTGTAT
301 GATTCTGTCA AACAGTTCTA CACCAAGGGC TCTGAGCATG CCAGCATTGG
351 GAGCCGCCCTC CTAGCAGGCA GCACCCACAGG TGCCCTGGCT GTGGCTGTGG
401 CCCAGCCCCAC GGATGTGGTA AAGGTCCGAT TCCAAAGCTCA GGCCCGGGCT
451 GGAGGTGGTC GGAGATACCA AAGCACCGTC AATGCCCTACA AGACCATTGC
501 CCGAGAGGAA GGGTTCCGGG GCCTCTGGAA AGGACCTCT CCAATGTTG
551 CTCGTAATGC CATTGTCAAC TGTGCTGAGC TGGTGACCTA TGACCTCATC
601 AAGGATGCCC TCCTGAAAGC CAACCTCATG ACAGATGACC TCCCTTGCCA
651 CTTCAATTCT GCCTTTGGGG CAGGCTTCTG CACCACATGC ATCGCCTCCC
701 CTGTAGACGT GGTCGAAGACG AGATACATGA ACTCTGCCCT GGGCCAGTAC
751 AGTAGCGCTG GCCACTGTGC CTTACCATG CTCCAGAAGG AGGGCCCCCG
801 AGCCTTCTAC AAAGGGTTCA TGCCCTCCTT TCTCCGCTTG GGTTCCTGGA
851 ACGTGGTGAT GTTCGTCAAC TATGAGCAGC TGAACGAGC CCTCATGSGCT
901 GCCTGCACCT CCCGAGAGGC TCCCTTCTGA
  
```





Fig. 1C.

restriction sites: fragment size indicated (Kb) when



**Fig. 1D.**

Partial sequence of human UCP2 gene , derived from hUCP2-g1 clone

.....

pSU04

.....

GAGCTCGAGGATCAACTAATGGTGACTGCTTGGCATCTCAGCCCCAGCCCCAGTCTCAT  
GGCATCCTCTGCCATACTGGAATCATAAGTTCTTGGTTGTCATGGCAACAGGATGGGAGT  
GGGAGTGGCTCAGGGCAGGCTCTCCCCATTGCTGGAGGCTCTTCCCATTGCTAGAGGATT  
GAGCCCAAGTACAAATCCCATAGAGCATCTTTCACGTGTGCAAAGGTCACCTGAGTGTTT  
ATGAAATCTGAGCAAATCTCGACTGTACTCATGGAAGTCCTAAGCAGACCTCCCCACAC  
CTGATATGTTGCTATAGGTGCTGAACGAGGATGGGTACTCCTCCACCATCAAGGACAAGA  
TCCTGACCATTGATGTGAACCCGGTTGGAGGCAGGGCACACGCATCACCTTTGAGAAGG  
AAGGGGACCAGGTGAGGGGGGAAGAAGCTGACTCAGGTCAGTCACTGAGCTC

.....

pSUB12sp6

.....

GGATCCCTAGCTGGGGCAGGGGCAGGTGGAGGGGGGTTGTGGTGGTGAGACTCCAAATGG  
AAAATATCCTGGCCAGGCACAGTGGCTCATGCCTGTAATCCCAGCACTTTGGGAGGCTGA  
GGTAGGTGGATCACTTGAGGTCAGGAGTTTGAGACCAGCCTGGCCAACATGGCAAAACCC  
CGTTTCTACCAAAAACACAAAATTAGCCAGGTATGATGGTGTATGCCTGTAATCCCAGC  
TACTTGGGAGGCTGAGGCAGGAGAATCGCCTGAACCTGGGAGAGGTTGCAGTGAGCCAAG  
ATTGCACCACTGCACTCCAGCCTGGGTGACAGCAAGTGAGACTCTGTCTCAAACAACAA  
AAAACACNAATGGAAAGTATCCTCAGGCAGAGACAAATGTTTGCCCTGCAGCCTCCACCA  
CCTCTGTGATTGAGGCCGCTCCCCCATCA

.....

pSUB14sp6

.....

GGATCCCAAGGAGGCGCAAGAAGCCAGGGCCTTGGCTGCACAGGGGTCTCCGCTTCTCTG  
TCCCTGTTCTTGTACCTGCACTCAGAGGCAGGTGGGCAGGGGTACTACAATTTCAAGGA  
GTGGAGACTGTGAGGTCCTGGAATCCCAAGGCATCTCCTATAGGGCTGGGCCCTTAGAAT  
TATGTCACTCAGACCCAGTTTGTAGGTGTCTGAAGAACTGAGGCCTGACACAGGTGATG  
CAGGCAAGAACACCCAGAAAGTCCACTACTGAACTGGGACCGGGACCCAGTCCTCCTTCC  
CCTTGTGGACTCCCCAGAGACCAGTGCTGGGGTCTTGGGGAAGCCTGTTTGGCAGCTG  
TGGAGCTAGGCCCTGAGAATACAACACCTCCCTCTTCCCTCAGCCTCAACCGCTGAACCA  
CTGCTGCTTCGCCCCCTGTAGCCCATGGTCAAACCTGGAGGCTAAACCTTCATGCTTGGT  
AAGGCAGGTGTAAATGGTCT

.....

pSUB14t7

.....

GGATCCCCAGTGTGCTACTAGCCAGCTGTGTGCCTTTGGGCAAATCACTTCACCTCTCTG  
AACCCGTTTCTTAAAGCTTACAGCCCCTACCTTGCAGGCCTTTCATGGAAACCAGATGGT  
GAGGCATGAAATGCTCTGGGGCAGGACTCAGGTGGAGTGGGATGGTGAGAGCTGCCCTCT  
TCTGAGGGCAANCCCTGAATTGGAGTGCANAGGCAGGGCTGTGAAATCTTCAACTCTAG  
GTCTCCTGTAGGGCAGCAGGAAAGCCTACTCCTTATCTTGAAGGAAGCCTGGGTCTTGGG

.....

pSUB23sp6

.....

GGATCCCTAATGTTCTTCCAAGGAACAGAGCCTTAGCCTTACAGTGCTTCTGTGAACAG  
GCAGCCCAGGGACTGTCCCCATTTACAGAGTGGGAAATGGGGGCCAGAGAAGAAAGGG  
ACTGCCCAGGGCAAGAACATGGGAGCTGAAAGGAGATCCCAGTCTGAACTCCAGTGTTT  
TCCTGATTTTCTTCAGTGCTTCTCAGAATAAAGCCCAATTACAGGAGCTACTCAGAAACC  
ATTTAATCATAATAATTTTATTTATTTATTTATTTATTTATTTATTTATTTATTTATTTGA  
GATGGAGTCTCACTGTCGCCCAGGCTGGAGTGAGGTGGCATGATTTGAGCTCATTGCAGC  
CTCCACCTCCTGGATTCAAGCAATTATCCTGCCCCACCTCCTGAGTAGCTGGGANTACAG

**SUBSTITUTE SHEET (RULE 26)**

**Fig. 1D. (continued 1)**

.....

pSUB23t7

.....

GGATCCCTAAGGAGAAGAGCCTGGAGGAGATGAGAAGTGGGGAGGAAGGCCTGCTGCGCA  
GATCTGGTGTACAGATCCAGGCGGTGCAGATGAGCAGATGGGACCAAGTGGATGGTGATT  
TAGAGGGTGGAAATCCACAGGGCTTAGTGACTGATGGCAGGTGGTTGTGGGAGGGGAGGCC  
AAGTCTAGGCCGGGGCAGGCAGGGCTCACTCTGATCTCCTCCCCGCTGCTCCTGCACTTT  
GCCCACTCACCTTGCCAAGAAGGATNGGGTTCACAAAAAAAGTTGTCCTTCNCCCCTGC

.....

pSUB5t7

.....

GGATCCCCCTAAGTTGGAGGAAAGGAACTGGCCCAAACAAAAAGGAGAGCAGTTTTCTC  
TGCATCAATGGTAGGCCAGGAGGAGTCTAATGCCCCAGAGTTTACTCTCAGCCCCCAA  
ATCACCTAGCTAAATGTTACCTTATCTAAGAAGTCCTTAGGTTTTTGGGGTTTGT  
TTTTTTTTGAGACAAGGTCTCACTCTCTCAGGAGAGTGGAGCAGAGTGGCACAATCAC  
AGCTCACTGCAGCCTCAACCTCCTGGGCTCAAGCAATCGTCCCAAGTAGCTGGGACTATA  
GGCCTGCACCACCATGTTGAGCTAATTTATTTTATTTTAAAAAGGTCTCATTAT  
TTGCCTGCTGGTCTTGAATCCTGGTTCAACATCTCCCACC

.....

ex12

.....

GAGCTCACATCGGTTTTCCCTCATGAGGCCACTTGGAGTCTTGCTGAGGGACTTGGTTCT  
ATTAGGGAAGGTGAGTTTGGGGATGGTGAGCAGGGAGGGCCTGGGGACATTGTGGCTAAT  
GGGGCTTTTCTCCTCTTGGCTTAGATTCCGGCAGAGTTCCTCTATCTCGTCTTGTTGCTG  
ATTAAAGGTGCCCTGTCTCCAGTTTTTCTCCATCTCCTGGGACGTAGCAGGAAATCAGC  
ATCATGGTTGGGTTCAAGGCCACAGATGTGCCCCCTACTGCCACTGTGAAGTTTCTTGGG  
GCTGGCACAGCTGCCTGCATCGCAGATCTCATCACCTTTCCTCTGGATACTGCTAAAGTC  
CGGTTACAGGTGAGGGGATGAAGCCTGGGARTCTTGATGGTGTCTACTCTGTTCCCTCCC  
CAAAGACACAGACCCCTCAAGGGCCAGTGTTTGGAGCATCGAGATGACTGGAGGTGGGAA  
GGGCAACATGCTTATCCCTGTAGCTACCCTGTCTTGGCCTTGACAGATCCAAGGAGAAAGT  
CAGGGGCCAGTGCGCGCTACAGCCAGCGCCAGTACCGCGGTGTGATGGGCACCATTCTG  
ACCATGGTGCCTACTGAGGGCCCCGAAGCCTCTACAATGGGCTGGTTGCCGGCCTGCAG  
CGCCAAATGAGCTTTGCCTCTGTCCGCATCGGCCTGTATGATTCTGTCAAACAGTTCTAC  
ACCAAGGGCTCTGAGCGTGAGTATGGAGCAAGGGTGTAGGCCCTTGGCCCTTTTTTCTC  
AGTGATGATTGATCTTAGTTCATTAGCCATATAGTTTTTATGGCCCCACGATCCCTAGG  
AAGATCAGGGGGAACAGAGAACTGGAAGGGGCCCTGGTCCTCCACATAGTTCCTAAGCAC  
CTGGGGCTATACCAGGCTCTGAGCAGGGCGTCATCCCATCACAGTCTTCAACACCACCTTG  
GGAGTAGGTAGTATCATCCAGTGTTATAGAAGAAGAGACTGAGGTGGGAAGGCAGTGGG  
TAGAGTGGGGACTTGGCCAGGGGACACAGTAGAGAGCCAGAAAAACACACAGTAGAGAGC  
CAGGACACTCGTCTCTAAGGCCAGCGTTCTTCCCTTTCACCTCCTTAGTATGCCATGCCA  
ACCCTCCATTTTACACATGACGAAACAGAGCCCCAAACAAAAGGTTGTCTTTCCAGATC  
ACATGGCAGGAAGAAGTAAAGCTGACCTGAGATCCCAAGTCTTAGGAATCCCAGTCCTCA  
GAAAGCCACTTCTCTCTGAGCCTTGGTTTTTACATTTGTGAGATGGAATGATTGTGATT  
TCTCAGGGCTGTTGAGCAGGTAAATGAAAATGTTTTATGAAAGAAAGCAAGTTTCAT  
TTTGGTCTTAGCCCTTGCTATGTCCCTAGCAAGAAGTAGATATTCATAGGGATATTTGT  
TTGATGTGAGGAGTCTTACAGCAAGAGCTTGTAGAAGGCCAAAAGCTTCTGGATTCTAA  
TCCAAAAGCAGGAGATGACAGTGACAGGGTGGTTTTTGGTGAGGAGAGATGAGGTAGAAA  
ATGAGTGCAAGCCCGCTGGCCACTGACCCCATGGCTCGCCACAGATGCCAGCATTGGGG  
CNTNCCNTTGG

**Fig. 1D. (continued 2)**

.....

ex34

.....

CCCGNTGNGCTGTGGCTGTGGCCCAGCCCACGGATGTGGTAAAGGTCCGATTCCAAGCTC  
AGGCCCGGGCTGGAGGTGGTCGGAGATACCAAAGCACCGTCAATGCCTACAAGACCATTG  
CCCGAGAGGAAGGGTTCGGGGGCCTCTGGAAAGGTGTGTACCAGTTGTTTTCCCTTCCCC  
TTTTCTCCTCCCCGATACTCTGGTCTCACCCAGGATCTTCTCCTCCTACAGGGACCTC  
TCCCAATGTTGCTCGTAATGCCATTGTCAACTGTGCTGAGCTGGTGACCTATGACCTCAT  
CAAGGATGCCCTCCTGAAAGCCAACCTCATGACAGGTGAGTCATGAGGTAGACGGTGCTG  
GGTCTCACCCCTTCCCCCATGCCAGGARCAAGGTGCGGGGGTCTAGCTGACACCAGAAGACC  
ACATCTTTTCATCCTATTTGCCCTTTCAGGGAGAGTAAGATATCTTACTTGCCATAT  
TGAAGCCAATTGGGATGAAGCTCCCACTTTCACATTGAGGAACTGAGGCTAGATTGGCA  
AAATGACTCTTTCAGGTCTCAGAAGATGTCTCAGCTGGAGTCCCTGTCTGTTTTGTTT  
TTTTGTTTGTGTTTTTTTTTTTTTTTTGAGATAGANTCTCACTCTGTTACCCGTGTAAT  
CTCAGCTCACTGCAACCTTCTCCTCCTGGGTTCAAGCAATCCTGTGCCTCACCTCCC

.....

ex56

.....

GTCATGGAGCTTGACTTCCGGATTAAGCATCTTCCACTTCAGCCTCTNGAGTAGCTGGGC  
TGA CTACAGGCATGCACACTGTGCCTGGCTAATTTTNGTATTTTGTAGAGACAGGGTTTT  
TGCCATGTTACCCAGTCTGGTCTTGA ACTCCTGGGCTCAAGTGATCCACCCACNTCGGCC  
TCCAAAAGAAGTCTGGATTACAGGCATGAGACATTGTGCCCAGCCTCTCTGTCTCTTTA  
AAATGAGAAACTCGTAGCTACTTAAGTAATTCCTGCCTTCTGGAATGATGGGTGAA  
GATCTTGACTGCCTTGCCTGCTCCTTGGCAGATGACCTCCCTTGCCACTTCACTTCT  
GCCTTTGGGGCAGGCTTCTGCACCACTGTCTATCGCCTCCCTGTAGACGTGGTCAAGACG  
AGATACATGAACTCTGCCCTGGGCCAGTACAGTAGCGCTGGCCACTGTGCCCTTACCATG  
CTCCAGAAGGAGGGCCCCGAGCCTTCTACAAAGGGTGAGCCTCTGGTCTCCCCACCCAG  
TTCAGGCNTCTTGGCTATGCATGTCTATTATGGGTGGGAGAGAACCACCTGGAAGTGAGT  
AGCAGCCAAGTGTGACTATTTCTGATCCTGGTCGTGGCATTTCACCAGCATTACCTATC  
CCCTTAATTCTTCCCTCCCAGAATTGCTACCATCACTGTTTATTAGGTGTTAAATGGAGA  
CTCAAAGGGAATTCATGCTTATAGCCAAGCAGCTGTGAGCTCAGTTCATTGAGTCTCCC  
AGCCTCCTTTGGGACAGAGCAACTGGGTTGGATTGAATACCAGGCCCAAGTGAGGGAAGTG  
GGAGGTGGAGGTGCCCCCATGACCTGTGATTTTTCTCCTCTAGGTTTCATGCCCTCCTTTC  
TCCGCTTGGGTTCTTGGAACGTGGTGATGTTCTGTCACCTATGAGCAGCTGAAACGAGCCC  
TCATGGCTGCCTGCACTTCCCAGAGAGGCTCCCTTCTGAGCCTCTCCTGCTGCTGACCTGA  
TCACCTCTGGCTTTGTCTCTAGCCGGGCCATGCTTTCCTTTCTCCTTCTTCTCTTCC  
CTCCTTCCCTTCTCTCCTTCCCTCTTCCCCACCTCTTCTTCCGCTCCTTTACCTACCA  
CCTTCCCTCTTCTACATTCTCATCTACTCATTGTCTCAGTGCTGGTGGAGTTGACATTT  
GACAGTGTGGGAGGCCTCGTACCAGCCAGGATCCCAAGCGTCCCGTCCCTTGGAAGTTTC  
AGCCAGAATCTTCGTCTGCCCCGACAGCCCAGCCTAGCCCACTTGTCATCCATAAAGC  
AAGCTCAACCTTGGCGTCTCCTCCCTCTCTTGTAGCTCTTACCAGAGGTCTTGGTCCAAT  
GGCCTTTTTGGTACCTGGTGGGCAGGGGAGGAACCACTGACTTTGAAAATGGGTGTGAT  
CCACCTTCCACCTCCAGCATCCAATCTGAAGCCCGTGTAGGTCATCTGGTCCATTTCTCT  
CTAGATCCAGGCCCTGTTACTAACATGGGGAGTGCAGGAGCCACCTGAGAGACAGCAGTG  
CCTCCCCTTCTTTGCCGGGCCACTTGAGCTC

**Fig. 1E.**

LOCUS hUCP2.genebank 1612 bp  
 DEFINITION hUCP2.genebank, 1612 bases.

```

1  TGA CTGTCCA CGCTCGCCCG GCTCGTCCGA CGCGCCCTCC GCCAGCCGAC
51 AACACAGCCG CACGCACTGC CGTGTTCTCC CTGCGGCTCG GACACATAGT
101 ATGACCATTA GGTGTTCTGC TCCCACCCAT TTTCTATGGA AAACCAAGGG
151 GATCGGGCCA TGATAACCAC TGGCAGCTTT GAAGAACGGG ACACCTTTAG
201 AGAAGCTTGA TCTTGGAGGC CTCACCGTGA GACCTTACAA AGCCGGATTG
251 CGGCGAGTTC CTCTATCTCG TCTTGTGCT GATTAAAGGT GCCCCTGTCT
301 CCAGTTTTTC TCCATCTCCT GGGACGTAGC AGGAAATCAG CATCATGGTT
351 GGGTTCAAGG CCACAGATGT GCCCCCTACT GCCACTGTGA AGTTTCTTGG
401 GGCTGGCACA GCTGCCTGCA TCGCAGATCT CATCACCTTT CCTCTGGATA
451 CTGCTAAAGT CCGGTTACAG ATCCAAGGAG AAAGTCAGGG GCCAGTGC GC
501 GCTACAGCCA GCGCCCAGTA CCGCGGTGTG ATGGGCACCA TTCTGACCAT
551 GGTGCGTACT GAGGGCCCCC GAAGCCTCTA CAATGGGCTG GTTGCCGGCC
601 TGCAGCGCCA AATGAGCTTT GCCTCTGTCC GCATCGGCCT GTATGATTCT
651 GTCAAACAGT TCTACACCAA GGGCTCTGAG CATGCCAGCA TTGGGAGCCG
701 CCTCCTAGCA GGCAGCACCA CAGGTGCCCT GGCTGTGGCT GTGGCCAGC
751 CCACGGATGT GGTAAAGGTC CGATTCCAAG CTCAGGCCCC GGCTGGAGGT
801 GGTGCGAGAT ACCAAAGCAC CGTCAATGCC TACAAGACCA TTGCCCCGAGA
851 GGAAGGGTTC CGGGGCCCTCT GGAAAGGGAC CTCTCCCAAT GTTGCTCGTA
901 ATGCCATTGT CAACTGTGCT GAGCTGGTGA CCTATGACCT CATCAAGGAT
951 GCCCTCCTGA AAGCCAACCT CATGACAGAT GACCTCCCTT GCCACTTCAT
1001 TTCTGCCTTT GGGGCAGGCT TCTGCACCAC TGTCATCGCC TCCCCTGTAG
1051 ACGTGGTCAA GACGAGATAC ATGAACTCTG CCCTGGGCCA GTACAGTAGC
1101 GCTGGCCACT GTGCCCTTAC CATGCTCCAG AAGGAGGGGC CCCGAGCCTT
1151 CTACAAAGGG TTCATGCCCT CTTTCTCCG CTTGGGTTCC TGGAACGTGG
1201 TGATGTTTCG CACCTATGAG CAGCTGAAAC GAGCCCTCAT GGCTGCCTGC
1251 ACTTCCCGAG AGGCTCCCTT CTGAGCCTCT CCTGCTGCTG ACCTGATCAC
1301 CTCTGGCTTT GTCTCTAGCC GGGCCATGCT TTCCTTTTCT TCCTTCTTTC
1351 TCTTCCCTCC TTCCCTTCTC TCCTTCCCTC TTTCCCCACT TCTTCCCTTC
1401 GGGCTCCTTA ACCTACCAAC CTTCCCTCTT TCAACATTCT CATCTACTCA
1451 TTGTCTCAGT GCTGGTGGAG TTGACATTG ACAGTGTGGG AGGCCTTGTA
1501 CCAGCCAGGA TCCAAGCGT CCCGTCCCTT GGAAAGTTCA GCCAGAATTT
1551 TTGTCTGCC CCCGACAGCC CAGCCTAGCC CATTGTCATC CATAAAGCAA
1601 GCTCAACCTT GG

```



## hUCP2.CDSF1

Forward Oligo: 5'&gt; CATCTCCTGG GACGTAGC &lt;3'

>> Length: 18  
>> Melting Temperature: 56.6  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 42  
>> % GC Content: 61  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> Longest palindrome has 4 bases (bases 12 to 15).

## hUCP2.CDSR1

Reverse Oligo: 5'&gt; GGAGAGAAGG GAAGGAGGG &lt;3'

>> Length: 19  
>> Melting Temperature: 60.1  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 57  
>> % GC Content: 63  
>> Hairpin Stem: 0  
>> Primer Dimer: 0  
>> No Palindromes.

&gt;&gt; Product Size: 1098

&gt;&gt; Annealing Temperature: 59.4

Try T = 51.6

## Pair 2

## hUCP2.CDSF2

Forward Oligo: 5'&gt; AGGTGCCCCCT GTCTCCAG &lt;3'

>> Length: 18  
>> Melting Temperature: 61.3  
>> Max AT Run Length: 1  
>> GC Clamp Strength: 32  
>> % GC Content: 67  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> No Palindromes.

**Fig. 1F.**

## hUCP2.CDSR2

**Fig. 1F. (continued 1)**

Reverse Oligo: 5'> TCAGGTCAGC AGCAGGAG <3'

>> Length: 18  
>> Melting Temperature: 58.7  
>> Max AT Run Length: 1  
>> GC Clamp Strength: 32  
>> % GC Content: 61  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> No Palindromes.

>> Product Size: 1043  
>> Annealing Temperature: 60.1                      Try T = 53.7

---

Pair 3  
hUCP2.CDSF3

Forward Oligo: 5'> CATCTCCTGG GACGTAGC <3'

>> Length: 18  
>> Melting Temperature: 56.6  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 42  
>> % GC Content: 61  
>> Hairpin Stem: 2  
>> Primer Dimer: 6  
>> Longest palindrome has 4 bases (bases 12 to 15).

hUCP2.CDSR3

Reverse Oligo: 5'> AGAGAAGGGA AGGAGGGAAG <3'

>> Length: 20  
>> Melting Temperature: 58.9  
>> Max AT Run Length: 2  
>> GC Clamp Strength: 47  
>> % GC Content: 55  
>> Hairpin Stem: 0  
>> Primer Dimer: 0  
>> No Palindromes.

**Fig. 1F. (continued 2)**

&gt;&gt; Product Size: 1096

&gt;&gt; Annealing Temperature: 59.4

Try T = 51.6

---

**Pair 4**

hUCP2.CDSF4

Forward Oligo: 5'&gt; CATCTCCTGG GACGTAGC &lt;3'

&gt;&gt; Length: 18

&gt;&gt; Melting Temperature: 56.6

&gt;&gt; Max AT Run Length: 2

&gt;&gt; GC Clamp Strength: 42

&gt;&gt; % GC Content: 61

&gt;&gt; Hairpin Stem: 2

&gt;&gt; Primer Dimer: 6

&gt;&gt; Longest palindrome has 4 bases (bases 12 to 15).

hUCP2.CDSR4

Reverse Oligo: 5'&gt; AAGGAGAGAA GGGAAGGAGG &lt;3'

&gt;&gt; Length: 20

&gt;&gt; Melting Temperature: 58.9

&gt;&gt; Max AT Run Length: 2

&gt;&gt; GC Clamp Strength: 43

&gt;&gt; % GC Content: 55

&gt;&gt; Hairpin Stem: 0

&gt;&gt; Primer Dimer: 0

&gt;&gt; No Palindromes.

&gt;&gt; Product Size: 1100

&gt;&gt; Annealing Temperature: 59.4

Try T = 51.6

---

**Pair 5**

hUCP2.CDSF5

Forward Oligo: 5'&gt; AGGTGCCCCT GTCTCCAG &lt;3'

&gt;&gt; Length: 18

SUBSTITUTE SHEET (RULE 26)

**Fig. 1F. (continued 3)**

```
>> Melting Temperature: 61.3
>> Max AT Run Length: 1
>> GC Clamp Strength: 32
>> % GC Content: 67
>> Hairpin Stem: 2
>> Primer Dimer: 6
>> No Palindromes.
```

## hUCP2.CDSR5


Reverse Oligo: 5'> AGGAGAGAAG GGAAGGAGG <3'

```
>> Length: 19
>> Melting Temperature: 57.4
>> Max AT Run Length: 2
>> GC Clamp Strength: 43
>> % GC Content: 58
>> Hairpin Stem: 0
>> Primer Dimer: 0
>> No Palindromes.
```

```
>> Product Size: 1125
>> Annealing Temperature: 60.8
```

Try  $T = 52.4$

Fig. 2.









**Fig. 3B.****Fig. 3C.****Fig. 4.**

Fig. 5.



**Fig. 6.**

HIB 1B cells grown in DMEM + 10% charcoal-stripped serum (*Media*) or with the addition of the thiazolidine dione BRL 49653 and the RXR $\alpha$  ligand LGD-1069 ( *TZD/RXR*). The results are the average of 2 independent experiments of 4 samples each. \*, significantly different from *Media* samples,  $p < 0.001$ .



## Map of the mouse UCP2 gene.

Clone "MMU2-L2" (insert 13.9 Kb) deposited at Pasteur Institute  
(Registration number: I-1868, 04.16.1997)




Fig. 7.

SUBSTITUTE SHEET (RULE 26)

**MMU2-L2: partial promoter sequence (unmapped)**

**SUBSTITUTE SHEET (RULE 26)**

**promoter and coding sequence:**

**SUBSTITUTE SHEET (RULE 26)**

Fig. 8B. (continued 1)


ATCTCAGGACCTCTCTTCGGCTGCCCTGGAGCTTTAGATGCAGACCGTGCCTGCACAGGATTATCTTTCTAGGCCCTGCGTT  
 TTAGTCATAGAAACCTTTAGTACAGGTGATCTTTGCTCTGGTCCCAACCTCGAGGACCCGACCTTTGGCAAAACCCCTAC  
 GCTGCATTACAAATGGCAGCTGAGCCACATGTTTTGATCTGGCAGCAACTGGCAGCTGAGCCACATGTTTTTGTATCTGG  
 CAGCAACTGAGACCTCTCTTTCTGGACTTCACAACATCTACCACTTAGCTCATGATtAGACAGGCCtGCaTCaGACtTC  
 aTGCaCTAAATACcTACAgAggAggTcTGCCCCagAgCAtCtCgAAcTgTTAAATGgTCAGAAAGGCTTCCATTTGTGAA  
 GGGCTTTATCTGAGCCCGAGACTGAGAAAACCGGGAGAcATGAGGCACCCGGGGCTGGGCAGACAGTGAGCCtGTGGGA  
 CATGGTTCTGAGCTCTCCCGTTTAAACGGGGACCCAGGGAGGGCAGGTGCCCTTCCAGGCTGCTGTcAGGATTTCTcAGTGG  
 AGTCGGAGTGGGAGCACTACCTGGGAGAGACTCCCCCGTGAGGAGTCAAGGAGCACCTTACCTAGACATTTGACTCCT  
 ATTCTCCCTGAGCTCCAGGGCCCCAGACTCTGACTGAAGTCAAGCTTTTCCCTAGAGGCGCCCTTTGAGCCCTGTTGTCTCATG  
 GTGAGTCCCTCTGATGTCTCTGTTTGTCCAGCCCTGGGTGAGCTGTTTGTCTTCCATGGCTGTCTGTGTTCACAGSTGAC  
 TACCGGCCACATCCTCCCTCCACCCCGTAGGGCTTTCTCTCTCTGCCCCAGCAACACAGGCAGCCCAATCAAGATGGGCTCA  
 GAGCGATTGCTCCAATGAAGCGGAGAAAGTGAACTAGGGAAGAAGAAAGAGTTCACCTCCCCACCCcACCCcAC  
 CCCACCCCGCTGdGTCTGGGCTCACTTCTCTCTAGGCCCTTAGAGGAgGGTGGTGTTCAGCTCAAGAACAAAATGGA  
 CCTTTTGGTTTCTCTGTATCTTTCTGCAATAGCTGTGAGCTGTGCCCTCCCATCGAACCCACAGATAACTGAgCATTTCCCGGA  
 CACTGGAGGAAAACAAACAGGCAACCTGGCTGGCCAGGTGCCCTTGCCcAGTGCCCTGGCCATGAGCCtGCAGGGGTAGGCA  
 GGCTCCATCTGTAAACCTGAATGGGGCCGGAGCTGGTGGGGCTGTAGGATGGAATTCATCAAGGTGTCTCATGTCTC  
 CCTTCATAATTACATACCAGACAGGACCTCTCTCTGTTTAGGATCCAGAGCAAAAGACTGAGGGgTTTGCCCTGCCaGtTC  
 CTGCTCTGATAGTCTTTGTCTTAGTCCAGCCGTTGGGATAAGTCCACAGATCAGGAACGGGGCTTCTCTCTGCAGTCTg  
 CCAGGAGATCAGAgGcTgTtGgtGgtTcGAgGGtGacCTGAGCCACACCCcCTATACCAATAAAGAGAGGAgTtCTT  
 CActGAatCTCAGTtCCCTGGCAGGAATtGCTtCCcAGACCCTActGAcTATTTTGGAGCTCTCTCTCTGGTTTcATCC  
 TGGACCCCTCTGAGATGTGCTTgTgCTTGGGgAAGGCTGGGCAGGGgTAGGTAGATGGTgGCCAGGgAgGGGCTGCT  
 GATACTGATGGCTTCTCCCTCTTGGCTTAGAacTCCGGCAGAgTCCCTtGCTCTTGGCGATtGAAGgTCCCGT  
 TTCTCCAAATtCTCTCCaTCTCTGGgAgGTAGCAGGAaTCAGAAATCAGTtGTTtCAAGGCCACAGATGTgCCCC  
 AACAGCCACTGTgAAGTTCTTGGGGCTGGGACAGCTGCCTGCAATGCAGATCTCATCTCTCTCTCTGATACCCGCA  
 AgTCCCGCTGCAGGTGAGATGGAgTGAAGTCTCTGATGTGTCTAATCTGCTCCCTCTCTCTGGGTCAgACTCTTGGCA  
 GGGATgAgGAgCTTGGAgCTGGCGGAGATCAGCAGAgCCATCTCTCTGCTGCACAAACCACCCCATCTTGGTCTTTTCAG  
 ATCCAAGGGGAGAGTCAAGGGCTAgTGGCACCCGACCCAGCGCCCAgTACCGTGGCGTTCTGGGTACCATCTTAACCAT  
 GGTGGCACTGAGGGTCCACGAgCTCTACAAATGGGCTGGTCTgCCGGCTGCAGCGCCAGATGAGCTTTGCCCTCCGTC  
 GCATTGGCCCTCAcGAcTCTGTCAACAGTTCTACACCcAAGGGCTCAGAGCGTGAATGCGGCAAGGACAGACTGACTG  
 ACCTTAGTCTGTCTGTGCAGCTCAGTGGCTCTTTTAgACTTTTCCaGaaACCCCGGaAGATcAGAGgTgGAcCTGATtT  
 CTGTATCTGAGCACCAGGGCTgTCTCAAGCAGGATGTgAGCCCATGTATgTCTTTaAAACCATCACGGTGTGAGCTgG  
 CCTcCTGGGTgACaCTtGGATgAggCTTGAATAGGGGAGGGACTGGCCAGAGCACAAATGGAGAGGCCAGCCACCA  
 CCTCTAGCCCCCACAACCCACATCTTCTTCTTACTATGCCCTCTCTAGAGGGGAGGGGGGTCTgAgACAGGAGTG  
 TCACCTTGACAGCTCTGGCTGGCTGGTGTTCGAACAGCTCTCTCTCTCTCTAGCCCTTCCAAATGCAGgCATtaAGtGtGtG

Fig. 8B. (continued 2)

TcACAAAGCCCTCCAGGATCGCTcTcCAGGGAGAAGtAGAGCTGGGgCTAAGAAATcTGCCcTcTgGACtCAGcTcTGcT  
 AGgAAAGccTcTcTgCTGgTcTtGgTtTcACATtTGATAAATGgAGcTgACCCcTGAATtACCAAGGCTGCCGAGTAG  
 CGTAAATGAACACATtTATGTGAGAAAGcGAGTCTCCATtTTGTGtTACATCCtGGTGTAAACCCtGCAAGAGTGATCT  
 AGAGCCTCTCTCTTCCCGTGGCATGGTGGTCTGGCAGAGCATGATGCACtTGAGTGAAGCCTGCGGGCCACTAACCCC  
 GCCCCCACCCcTCTTGCCACAGATGCAGGCATCGGGAGCCGCCCTCCtGGCAGGTAGCACACAGGTgCCCTGGCCGTGG  
 CTGTAGCCACGCTACAGATGTGTTAAAGGTCCGCTTCCAGGCTCAGGCCCGGGCTCTGGAAAGGTGTGTGCTGGGTGTGGCTTT  
 ACTGTGGAAGCCtACAAGACCATtGCACGAGAGGAAGGATCCGGGGCTCTGGAAAGGTGTGTGCTGGGTGTGGCTTT  
 TGTCCCTCTCTCTCCATGCCCTGGCTGACCTACACAGGTCTCTTTTCTACAGGACTTCTCCCAATGTGTGCCCTAATG  
 CCATtGTCAACTGTGCTGAGCTGGTGACCTATGACCTCATCAAGATACTCTCTGAAAGCCAACTCATGACAGGTGTG  
 TTAGCGCCACTTCAAGCTGGTAGGGGCGTCTAGTGTCTAAAGACTGCATctTGTtATCTCTGCTTGCCTCGTGGTtTTT  
 AGAGTTTATATTTCTTACCATTCAATCCAAACAGGGTAAACCCCACTTTGCACATAAAAGCCCTCTGCGGCAATCAACAAAG  
 ACTCTGAAGTCCCCtAAGAGAGCTAGCTGTAGTCTGTCTTTAAACAAATGAGACcCTCTGCGGCAATGGTGGGAtGC  
 GgCTgGGAAGcTgAcAgCcTcTcCCCTGCTTCATGSCAGatGAcCTCCctTGcCACTTCACTcTGCCtTcTgGGCcg  
 GcttctGCACCCGTCATCGCCCTcCCCTGTGTGAtGTGtTCAAGACgAGATACATGAACtCTGCGCTTGGGCCAGTACCAC  
 AGCGAGTCAcTGTGCCCTTACCATGCTCCGGAAGGAGGCCCGcGcCTTCTACaAGGgTGAGCCctCagATcCCCT  
 CaAgAAcTCCAGAGAAGTGAGATGCAGCCAAATGTCAAGACGCCCATAGTtGCTGTACcTCTGTtTTtTAGAAGTTAA  
 AATGAGCAATTCAGAcTGGGGTtGGGATGGGTGGGTCTAGTGGTGGAGGGCTTGTGTAGCATGCTTGGAGACCCtGGGTT  
 CTGCTCCCAAGCAGCAGGAGTAGGGAGCTGGGTACTCGGGATtCTTGGGCATGTGGCCATGGGTTGGGTGGATGGATG  
 AGCCAGTGGTtAGAAcGAGGTGGTGGAGGGACCTCCATGACTGTGTtTGTCTCTCTAGGTTCATGSCCTTCTCTCTC  
 CGCTGGGATCCtGGAAcGCTAGTATGTtTGTACCTATGAGCAGCTCAAAAGAGCCCTAAATGGCTGSCCTACCAATCTCG  
 GGAGCCACCTTTCTGAGCCTCTCCATGCTGACCTGGACCCCTGCTTCCAGCCCTGCTCTCTTCTTCTCATCTCTCTGC  
 CCAGTCCCATTTCTTCCCATTTCTCTGACCCCGATTTACTTCCACCTCACCTCCCTGTGCTCTGTACTGATGACTCA  
 CAGTAGGAGGCCtGACACAGCCCTGAGCCCTCAGCCCTTTCTACAGCTAAGCCCACTTTCATCTTCATCTCCCCAGC  
 CCAGCCAGCCAGCTCAGCCAGCCCTTCAcCCATAAAGCAAGCTCAATGTtGGTGTCTTCTTCTCATATGTtTACAGAG  
 GTCTTGTtCCAAAGGAcCTTTTCAGGACcTTGGCAGGcAGGTGAGGACCCcAgTGTCTCTTAAAGAGGgAtGTGACCCACA  
 TtCTGCTGTAAACATCTGAGAGgGCTGGAGNGTCTtCCAGTCTGTCTcTCCAgACCCAgTGTGGCTCAGAgATCAGT  
 GGCACAGCCAGGGAGcACTGCAGcCTTTGTCCCTCTTACCCTtTCCGTGAACACCATGTAAAGCTGTCACtCAGTACCTGGT  
 ACTCCACCCACACTGGCCAGGCCCTGTCTATCTGCAGAGTGGAGTATCCCThCCTCCCACTCAACTTGTtTAGGAGA  
 TGTAGGGAAACTCCAGAACCTTCCATTTTGGGGGGGAGCTTACTTACTCTGTtTTACACATAGCAACCCACTCTCTGA  
 GGCCATAGACCTACAGAGACAAAGCAGGCTGGCCCTCAGCTATAATCAcACCCCTGAGACATGCTCTGAACCCGGGGTGGG  
 TGGGACAGGGCAGGtCTGACCCCATTTGAAGGGGCTCAGGCTCCCTAnCCAAAGATCTAGCCTGTAGGGGTGTGTGTGTG  
 TTTCAACnGgNcTTcACACTAGCACACTGGCCCTGGAACCTCGCTGTGTAGTCTAGAACTCATGACAAAtCTGCTGTAGTT  
 TCTCAAGGCTGGGATtACAGGCCACCTGTAGCACCCGACCTCTGAAGAGCCCTTCTGATGGAGCCCTTACCTTTTCCCCCA  
 AGACTGGGGAnCAGTCAACGCTTAGATTGTCAgGGTGTtTTCAGTCACTCTCTTCTCCCGCCAGCTCTCTCTCAgCCAGgAG



# MAP OF THE HUMAN UCP2 GENE clone hUCP2-g2



\* 3,8 kb Bam HI fragment containing exon1(sequence 2), intron 1, exon 2 (sequence 3) and a part of intron 2.

\*\* the 5' region upstream the +1 site encompasses 3kb DNA or more.

**MAP OF THE HUMAN UCP2 GENE  
clone hUCP2-g2**

**Fig. 10A.**

sequence 1

GGATCCATCCATCTTTCCTGCCAGTGCAGAAAATTGCATGGGTCATTCAC  
CTATTAGTTATTGAACAGCCTCGTCTCCCTTGTGCCCAACTCACAGTCTG  
GCACTGGACAGGTTCCTAAGGTGCCTTCTAAATAGGAGAGCCATGGTGAC  
CCTCCTCAGCCCCACCTTTGGCCCTGACCCTCCCTAGGCTCTCACCTGCT  
GCACTGTGGAGGTGAGGACCCTAGATGACCGTCTGCTCAACATCCCCATC  
AATGACATCATCCAGTGAGTCCATCTGCCTTGGGCCCCAGTAGCCAAGAG  
AAGGGCTAGCCTGGGATTTAGGTGGTAGTGTGACAGGTCTGCCCAGTAGA  
GTTGTACAGGCTGCCCCCAGCAGAAGGGTTCCCTGCTGAGGGGATaGAGA  
gGACCAAAACTGAACCcACACTGGCCAAGCTGTGGGCTGCAATCACCTGG  
CAAAAGGAacATCTTTTCTAATCCACACAaAGCCACTGgCAGAGgCTAA  
AGGCCCCATTGTTGGGCAACCTGTTACCTCCATGCCCTGCCAAACTTCTC  
ATAGAATTCTGAATTNGCTTATCCNAAGTTCCTTTTCCAGAAAGGGCCTC  
TGCCCGTGAAGAAGTTTGCTTGGCACCCACAC

**Fig. 10B.**

sequence 2

GANGAACCACCGGcGCGTTcGTTcGCAGGAGGTTGGTTAgTTTGCCcAGG  
GGTAAGGGGGGcTGGGCCCATAAAAGAGGAAGTGCAAcTTAAGACACGGC  
CCCcGTTGGACGcTGTtAGAAACCTTCcTGGGTTGGGAAGGCAAGAGGTG  
TGTGACTGGACAAGAATTGTTTcTGGGCGGTCAgTCTTGCCcATCcTcACA  
GAGGTTGGCGGCCCGAGAGAGTGTGAGGCAGAGCGGGGAGTGGCAAGGG  
AGTGACcATcTCGGGGAACGAAGGAGTAAACGCGGTGATGGGACGCACGG  
AAAACGGGAGTGGAGAAAGTCATGGAGAGAACCCTTAGGCGGGGCGGTCCC  
CGCGGAAAGGCGGCTGCTCCAGGGTCTCCGCACCCAAGTAGGAGcTGGCA  
GGCCCCGGCCCCGCCCCGCAGGCCCCACCCCGGGCCCCGCCCCGAGGCTT  
AAGCCGCGCCGCGCCTGCGCGGAGCCCCACTGCGAAGCCCAGCTGCGCG  
CGCCTTGGGATTGACTGTCCACGCTCGCCCGGCTCGTCCGACGCGCCGTC  
CGCCAGCCGACNGACACGGCCAAACGAAAGGNCCNNNNCCCTGCGgCt  
CGCGNACCCACGCTTGTCTGCGTGCGCTGCCCGCTCTTCCATTTACCT  
TCTCTCCACCCAAgTTTGTACTCTCTTCTCTCTCGGTGTtATATTTT  
GTGTTTGTGTGTGTGTgTGAGACAgGCGCTCGCTCTGTCTCCACGCTGG  
AGTGtCAGTGGCGGATATCgGCTCACTGCAGaCTCCACCTCCCAGGTGN  
AAACGATNCGCCTGCCGAGTATCTGGGATaACAgGCGCCCGCCACCACAC  
CCTGGcTAAAtATTTNTGTGTgTTTGTANAANATAGGGTgTTCGCCCACGT  
TgGtGcAGgCTGGNCTCTCAANTTGCTGAGATTCAAgCAATNTGCCCGCC  
CTCGGNGCTCACAAAAGTCNCTANAAANTTTAGGCGTGAAACCCCCCGG  
TNNNGGGCTGTTGCTAAANCNCCTCNTGTCCCTGGGGNCTCTAAAANCTN  
CTNCACTCCTCTTTCCTCAATCCCCTTGTCTTTTCCCCCNCCCGCTCAA  
TTNGNNGGTNNTTTGANNCCNCCTTTTNAATNCCCNCCCTTTNCANAAN  
NAAACTAACC



**Fig. 10C.**

sequence 3

ATTGGCCTTGGGNAGGGCCNGGTTCCCAGNCCTTTCCAAACTTTCTTNA  
CAGCCCGGACGNGGGGACNTAAGCCAATTTCCGGGGAGAGTTTNTGGTC  
CCANTGGNGACCCCTTNAAAAATTTCCNATTNACCAAAGTNTGATGG  
ACTGNGTTAGGGGGTGCTTATATAGAGTACTGAGTGTAAACAAAAGCAGA  
AGTCTGGATGAGAACCCAATTTGTGATATTAAGCAGGTGGGGTGGGGT  
GGGGAGTGTACCTAGGTTCAATTTCCGCCNTGCTTNTCCCCCTTTCCAGT  
GTGTGCACTTAACCAGTTCCTGGGCCCTGTTCCCCATCCCCCTCCAAG  
GCATGGATTGGGTGGGCTTGTGTGTCTTGGGGCAGGTGGCCCTTTCTAA  
ACTCTCTGCCTTTGCTCACCCACAGGACACATAGTATGACCATTAGGTG  
TTTCGTCTCCACCCATTTTCTATGGAAAACCAAGGGGATCGGGCCATG  
ATAGCCACTGGCAGCTTTGAAGAANGGGACACCTTTAGAGAAGCTTGAT  
CTTGGAGGCCTCACCGTGAGACCTTACAAAGCCGGGTAAAGAGTCCAGTC  
CAAGGAAGAGGTCTCTTGCTGCCTCCTAACCCCTGTGGTCTAGGGGCAGG  
AGTCAGCAGGGCATTAAACAAAATAATTACCATCCCCACCCCGACAGT  
GAAGTGGCTCTTTCCAGTTCACAGAGCACTCTCACACCTCCCCGCTCTC  
ATTCTGGCCCTTCAGCTGACTCGGACAAGCCAAGGATCTTGGTCCCCAT  
TTTATAAAGGAGAAAACCTGAGGCCNCGTGTNACAGTGATTGGCCCCAA  
GTCATCCCGGGAGCCAGCAGAAGAGCTAGGACAGGAACCTATTGTTCTA  
ACTTCATATTGATGCTAGCTTTTGACTATCCCTGAAACCGAAATTGGTT  
ATCAGCCCGGCTCTGAACTGGTTATTGCTGGGGACTGTTAAATAGGA  
TTAACTATTTCTATTCCTGCATTTTAATTGCTGTTAGTAGGGCAGCTTA  
CCACCCTCC

**Fig. 10D.**

sequence 4

TTCAGTGAAGCCTCGTGGGCAAGCTGGAGGGAAACAGGAATAAGTTCAGGC  
CCTGTTcTATAGGTCCCAGTGTAAGTTGCTATGGTGAGTATcTTCATTTCC  
CTGCTTGCCCCAGCCACcTGGAGTGAGAAGCCCAAGAGGAAGCTGGGTGA  
GCTGTTTGTTCATGGGTCTCTGTGTTTACAGCTGACTCCCTTCACCAG  
CCAGCCCTTTTACCTGAGCCCCAGCAACAAAGGCAGTCAGGCGGGGCTCA  
AAGCAGCTGCTCCAATGAAGTCAAAGAAATAAGCTCAGGGGAAGAAGCAG  
GTCACCCCTCCCCACTAGGGTGCTGGGCTCACTTCCTCCTGGGGCAGTGG  
AGGAGGGTGTGGTTCCAACCTCAGAACAAAATGGGGCTTTTGGTTTACTTT  
ATCACTCTTACAGCTCTGACCTGGACCCCTCATGCCCTGCCTGTCTTGT  
GGTGTAAGTGCGGATCCCCCTAAGTTGGAGGAAAGGAACTGGCCCCAAC  
AAAAAGGAGAGCAGTTTCTCTGCATCACATGGTAGGCCAGGAGGAGTCT  
AATGCCCCAGAGTTTACTCTCAGCCCCCAAATCACCTAGCTAAATGTTA  
CCTTATCTAAGAAGTCCTTAGGTTTTTTGGGGTTTGTTTTTTTTTTTTT  
GAGACAAGGTCTCACTCTCTCACCAGACTGGAGCACAGTGGCACAATCA  
CAGCTCACTGCAGCCTCAACCTCCTGGGCTCAAGCAATCGTCCCAAGTAG  
CTGGGACTATAGGCCTGCACCACCATGTCCAGCTAATTTATTTTTATTTA  
TATTTTTTAgaCAGGGTCTCATATATGTTGCCCTGGCTGGTCTTGAAGTCC  
TGGGTTCAAgCAGTCTCCACCTCTGCCTCCCAAAGTGCTAGGTTTTTTT  
TtGTTTGTGTTGTTGTTGTTTGTGTTTGTGAAACAGAGTcTTGcTCTGTGCGCc  
TAGGCTGGAGTGAGTGGCACgATCTCAGcTACTGCAACcTCCACcTCcT  
GGGTTCAAGTGATTcTCTGCCTCAGCCTCcTAAGTAGTTGGGAATACAG  
CGGTGTGCCAACACACCAGCTCATTTTTtGTATTTTTTAgaGGAATAGGGG

**Fig. 10D. (continued 1)**

TTTTGCCATGTTGGCCAAGCTGGTCTCAAACCTCCTGAcCTCAGGTGATTC  
GCCCCGCTCAGCCTCCCAAAGTGCTGGGTTTACAGGCGTGAGCCACCACA  
CCCAGCCCCAAGAAgTcTTTTcTGATcACCCAcTCTTcTTCTCTCCCAat  
GGCATtAgTtGTTCCCTCCTTTGCATTTTGAGAGTATGTCCtGTAAGCCC  
cAAATGcAgCtTGAATCATCTGCCCATCcACCCCCCTGTGCCCAACAGTAA  
GCCTCCTCTAGAGTAGATACTAtCTCCTGCATCTCAGTGAACCACTGCCc  
AgCAAAGCAGTCTTGCTAAAACAAtGACTcTAGAgATCCTAAGCTGTGtG  
AGAGCTGGAGGAGAGAATTAGAcTGATGgTCTGGGAAGGGATTGAATTAG  
TCATCTTGtACCTTTTCTTCTTGACTTAAGTTCCAGACCTGTAGCAACCA  
TTCCTGCTTAGACATCCAGAACATAAGCCTAtGGgtctGTGCCTGTTGGG  
TCTTAGTCTGGGTGAACTTTTCTCTACTTCTGTcAGCtCTCCAGATGAA  
CCACAGAAGCAGGAATGTGGGCATCATCAGTGAATCTCTGCATACAGCA  
GACAAAGGGCTGGTCCAGTGGCTGTTTATGAGGCAGCGCTAGGAGAGCTC  
TGATCCAGACTCTCCCTGCAGTGAAAGGGAGGGAGCCCTTCATGAAGTAT  
TGACTGCTTGAGCAGGAATTGCTTCACCAGCACCTAACTGAGTGCCTCTC  
GAGCTCACATCGGTTTTCCCTCATGAGGCCACTTGGAGTCTTGCTGAGGG  
ACTTGGTTCTATTAGGGAAGGTGAGTTTGGGGATGGTGAGCAGGGAGGGC  
CTGGGGACATTGTGGCTAATGGGGCTTTTCTCCTCTTGCTTAGATTCCG  
GCAGAGTTCCTCTATCTCGTCTTGTTGCTGATTAAAGGTGCCCTGTCTC  
CAGTTTTTCTCCATCTCCTGGGACGTAGCAGGAAATCAGCATCATGGTTG  
GGTTCAAGGCCACAGATGTGCCCCCTACTGCCACTGTGAAGTTTCTTGGG  
GCTGGCACAGCTGCCTGCATCGCAGATCTCATCACCTTTCCTCTGGATAC  
TGCTAAAGTCCGGTTACAGGTGAGGGGATGAAGCCTGGGAGTCTTGATGG  
TGTCTACTCTGTTCCCTCCCCAAAGACACAGACCCCTCAAGGGCCAGTGT  
TTGGAGCATCGAGATGACTGGAGGTGGGAAGGGCAACATGCTTATCCCTG  
TAGCTACCCTGTCTTGGCCTTGcAGATCCAAGGAGAAAGTCAGGGGCCAG  
TGCGCGCTACAGCCAGCGCCcAGTACCGCGGTGTGATGGGCACCATTCTG  
ACCATGGTGCCTACTGAGGGCCCCCGAAGCCTCTACAATGGGCTGGTTGC  
CGGCCTGCAGCGCCAAATGAGCTTTGCCTCTGTCCGCATCGGCCTGTATG  
ATtCTGTCAAaCAGTtCTACACCAAgGGCTCTGAgCgTGAgTATGgAgCA  
aGGGTGTAGGCCCCCTtGGCCCTTTTtCTCAGTGATGATtGATCTtAGTt  
CATtCagCCATATAgTtTtTtAGgCCCCAcgATcCcTAGgAAGatCAGGG  
GgAACAGAGAActGGAAGGGGCCCTGGTCCCTCCACATAGTTCCTAAGCAC  
CTGGGCTATACCAGGcTCTGAGCAGGgCGTCATCCCATCACAGTCTTCAA  
CACCACCTTGGGAGTAGGTAGTATCATCCCAGTGTtATAGAAgAAgAgAC  
TGAGGTGGGAAGGCAGtGGGTAgAGTGGGACTTGGCCAGGGGCACACAG  
TAGAGAGCCAGAAAACACACAGTAGAGAGCCAGGACACTCGTCTCTAAGG  
CCAGCGTTCTTCCCTTTTcAcCTCCTtAGtAtGCCAtGCCAACCcTCCATT  
TTACACATGACGAAACaGaGCCCCAAACAAAAGGTTGTCTTTCCcAGATC  
ACATGGCAGGAAGAAGTAAAGCTGAcCTGAGATCCCAAGTCTTAGGAATC  
CCAGTCCCTCAGAAAGCCACTTCTCTCTGAGCCTTGGTTTTcACATTTGTc  
AGATGGAAaATGATTGTGATTCTCAGGGCTGTTGAGCAGGTAAATGAAAA  
TGTTTTTATGAAAGAAAGCACCAAGTTTCATTTTGGTCTTAGCCCTTGCTA  
TGTCCTTAGCAAGAAGTAGATATTcATAGGGATATTTTGTtTGATGTGAG  
GAGTTCTTACAGCAAGAGCTTGTAAGAGGCCAAAAGCTTCTGGATTCTAA  
TCCCAAAAGCAGGAGATGACAGTGACAGGGTGGTTTTGGTGAGGAGAGAT  
GAGGTAGAAAATGAGTGCAAGCCCGCTGGCCACTGACCCCATGGCTCGCC  
CACAGATGCCAGCATTGGGAGCCGCCTCCTAGCAGGCAGCACACAGGTG  
CCCTGGCTGTGGCTGTGGCCAGCCACGGATGTGGTAAAGGTCCGATTC


**Fig. 10D. (continued 2)**

CAAGCTCAGGCCCGGGCTGGAGGTGGTCGGAGATACCAAAGCACCGTCAA  
TGCCTACAAGACCATTTGCCCGAgAGGAAGGGTTCCGGGGCCTCTGGAAAG  
GTGTGTACCAGTTGTTTTCCCTTCCCTTTTCTCCTCCCCGATACTCTG  
GTCTCAGCCAGGATCTTCTCCTCCTACAGGGACCTCTCCCAATGTTGCT  
CGTAATGCCATTGTCAACTGTGCTGAGcTGGTGACCTATGAcCTCATCAA  
GGATGCCCTCCTGAAAGCCAACCTCATGACAGGTGAGTCATGAgGTAgAc  
gGTGCTgGGTCTCACCCTTCCCCcATGcCaggAgCAGgTGCAGGGGgTcTA  
gCTGAcACcAgAagAccacATCTTTTCATCCTATTTGCCCTTTGCAGGGA  
GAGTAaGAtATCTCTTACTTGCCATATtGAAGCCAATTGGGATGAAGCTC  
CCACTTTGCACaTtGAGgAAcTGAgGCTAgAtTGGCAAAATGAcTCTTtC  
AGgTCCTCAGAAGATGTCTCAGCTGGAGTCCCTGtCtgTTTTTgTTTTT  
TgtttGtttGttTTTTGttTTTTtGaGAtAGaGTcTCAcTcTGtTACCC  
GtGTAATcTCAGCTCACTGCAACCTTCTCCTccTGGGTTCAAGcGATTCT  
TGTGCCTCAGCCTCCCgAgTAGcTGGGATGACAGGTGTGCACCAgCACAC  
tGGCTAATTTTTGTATTTTTtAGTAGAgAtGGAGTTTCACCATGTTAGCCA  
GGCTGGTctCGAAcTCCTGGCCTCaAGTGATCTGCCCCACCTTGCCCTCCC  
AATGTGCTgGGATTACAGGTGTgAGCCTcTGCGCCCCAtCcTcTTGtTtG  
tTtTTTGAgACAGGGTCTtGcTCGGTtGCCCCAgGctGGAgTGCAGTGGGG  
tgATTAATGGcTCATtGcAgCCTcgACCTCCctGAcTCAAgCAATCcTCC  
CACcTCaGCCTCcTGAgTAGcTGGGGCTGActAcaGGCaTgCACACTGTg  
CCTGGCTaATTTTTGTaTTTTGTAgAgACAGGGTTTTTGGCCATGTTACCC  
AgTctGGTCTTgAAcTcTGGGCTCaaGtGATCcaCCcAcCtcggCCTcc  
AAAAGAAGTCCTGGATTACAGGCATGAGACATTGTGCCCAGCCTCTCTGT  
CTCTTTAAAATCATGAAAACCTCGTAGCTACTTAAGTAATTCTCCTGCCTT  
CTGGAATGATGGGTGAAGATCTTGACTGCCTTGCCCTGCTCCTTGGCA  
GATGACCTCCCTTGCCACTTCACCTTCTGCCTTTGGGGCAGGCTTCTGCAC  
CACTGTcATCGCcTCCCCCTGTAGACGTGGTCAAGACGAGATACATGAAcT  
CTGCCCTGGGCCAGTACAGTAGCGCTGGCCACTGTGCCCTTACCATGCTC  
CAGAAGGAGGGgCCCCGAgCCTTctACaAAGGgTGAGcCTcTGgTCcTCC  
CcACCCAGTTcAGGCcTcTTGGcTATGCATGTCTATTATGGGTGGGAGAG  
AACCACCTGGAAGTGAGTAGCAGCCAAGTGTGACTATTTCTGATCCTGGT  
CGTGGCATTTTACCAGCATTCAcCTATCcCcTtAaTtCcTtCcTCCcagA  
aTtGcTAcCATCAcTGTtATTAGGTGTtAAATGGAGACTCaAaGGGAaT  
tCATGCTTATAGCCaAGCAGcTGTGAGCTCAGtTCAttGAgTCcTcCCAG  
CCTCCTTtgGGAcAGAgCAAcTGGGTtGGAtTGAATAcCAGGCCcAGTGA  
GGGAAGTGGGAGGTGGAGGTGCCCCCATGACCTGTGATTTTTCTCCTCTA  
GGTTCATGCCCTCCTTTCTCCGCTTGGGTTCTTGGAACGTGGTGTGTTT  
GTCACCTATGAGCAGCTGAAACGAGCCCTCATGGCTGCCTGCACTTCCCC  
AGAGGCTCCCTTCTGAGCCTCTCCTGCTGCTGACCTGATCACCTCTGGCT  
TTGTCTCTAGCCGGGCCATGCTTTCTCTTTCTTCTTCTTCTTCTTCTT  
CCTTCCCTTCTCTCCTTCCCTCTTTCCCCACCTCTTCTTCCGCTCCTTT  
ACCTACCACCTTCCCTCTTTCTACATTCTCATCTACTCATTGTCTCAGTG  
CTGGTGGAGTTGACATTTGACAGTGTGGGAGGCCTCGTACCAGCCAGGAT  
CCCAAGCGTCCCGTCCCTTGGAAAGTTCAGCCAGAATcTTcGTCCCTGCCC  
CCGACAGCCCAGCCTAGCCCAcTTGTcATCCATAAAGCAAGCTCAACCTT  
GGCGTCTCCTCCcTCTCTTGTAGCTCTTACCAGAGGTCTTGGTCCAATGG  
CCTTTTTTGGTACCTGGTGGGAGGGGAGGAACCACTGACTTTGAAAATG  
GGTGTGATCCACCTTCCACCTCCAGCATCCAATCTGAAGCCCGTGTAGGT

SUBSTITUTE SHEET (RULE 26)

**Fig. 10D. (continued 3)**

CATCTGGTCCATTTCTCTCTAGACCCAGGCCCTGTACTAACATGGGgAgT  
GCAGGAgCCACcTGAgAGACAGCAgTGCCTCCCCCTTCCTTTGCCGGGCCA  
CTTGAGCTCTTACTCAGAATCTGGTACTCTAGTGCCTGCCATCCCAACCC  
CCCACCCCAGCCGCAGGCCCTGTTTATCTGCACAACAAGAGTGCTCCTGTG  
TGCCCTGCATCTCCTGCAGTTCAGAGGAACATGAGACTCTTAGATGCTG  
TTGACTTTTATTTTATTCCATTTTACAAATGGAAGGAAGACCCACCTCCCC  
CAAAGTCCCAGACCTTGTGAGAACAAGTCAGCCTcCTTcCAcCCTCcACA  
GCCACAGCcAcACccACaGAGGAAATGTTACTGAACTGGGTGGAGCAGGC  
CctGAcTCcACAgAGGGTGGgTGGAGGCTGCAGGGCAAACATCTGGTCTC  
TGCCCTGAGGATACTTtCCATTtGTGTTTTTTGTTGTTTTGAGACAGAGTC  
TCACTTGCTGTCACCCAGGCTGGAGTGCAgTGGTGCAATCTTGgcTCACT  
GCAACCTCTCCCAGgTTCAGGCgAtTCTCcTGCCCTCAGCCTCCCAAGTAG  
CTGGGATTACAGGcaTACACCATCAtACCTGGCTAATTTTTGTGTTTtTG  
GTAGAAacgGgGTTtggccatggtggccaggctggtctcaaactcctgac  
ctcaagtgatccacctacctcagcctcccaaaGTGCTGGGATTACAgGCA  
TGAgCCACTGTGCCTGGccaggatattTTCCaTTtGGAGTCTCaCCacca  
cAACCCCCCTCCAcCtgCCCCTGCCCCAGCTAGGGATCCaaGGAGGccGC  
aAGAAGCCAGGGCCTTGgCtgAcAGGGGTCTCCgctTCTCTgTCCCTGT  
tCTTGTCACCTgcACTCAgAGGcAgGTGGGCAGGGGTACTACaaTTTCAA  
GGAGTGgAGACTGTGAGGTCCTGGAATCCCAAGGCATCTCCTATAGGGCT  
GGGCCCTTAGAATTATGTCACCTCAGACCCAGTTTGTAGGTGTCTGAAGAA  
ACTGAGGCCTGACACAGGTGATGCAGGCAAGAACACCCAGAAAGTCCACT  
ACTGAACTGGGACCGGGACCCAGTCCTCCTTCCCCTTGTGGACTCCCCCA  
GAGACCAGTGCTGGGGTCCTTGGGGAAGCCTGTTTGGCAGCTGTGGAGcT  
AGGCCCTGAGAATAcgAacACCTCCcTcTTCCCTcAGCCTcAAGCCGcTG  
AAGCCAcTGcTGcTTCGCCGCCCTcGTAAGCCCAATGgTCAGAGcTGGAGG  
cTAGACCCTTCAgTGCTTGGGTTGAGGGCcAgGGTGTTAgATTGGTTCTT  
GGAGAAGGAACGAGGGCCCCAgGATtCTtCAGCTTCTTAgtTTTTTGACAAA  
TTGAGCTGAGGCCCCaTAgtCcTCGGgAGGgACAGGGTTGAGTGCCATAA  
GTCGGCAAACCAgGGTAAAgGTGACAGGCAGCTCAgCCAGGCTGCAGGGG  
GTGGCATATACAgAGGACcTGGCCAcTACTTTATGTACCTTCTTACACTA  
ATTCTGTGAGGCAGGCTGTTTGTAgCTCTGCTCTGGACGGGAAGAAgTA  
GGGGCAgTTTGGTAGATGTGTGTCAAAG



**Fig. 11.**

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>16</u> , lines <u>21-25</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution  <p style="text-align: center;">American Type Culture Collection</p>	
Address of depositary institution (including postal code and country)  <p style="text-align: center;">12301 Parklawn Drive Rockville, MD 20852, USA</p>	
Date of deposit  <p style="text-align: center;">April 18, 1997</p>	Accession Number  <p style="text-align: center;">97993</p>
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<p style="text-align: center;">For receiving Office use only</p> <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;"> <input type="checkbox"/> This sheet was received with the international application       </div> <div style="border: 1px solid black; padding: 5px; min-height: 40px;">         Authorized officer       </div>	<p style="text-align: center;">For International Bureau use only</p> <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;"> <input checked="" type="checkbox"/> This sheet was received by the International Bureau on:          23 February 1998 (23.02.98)       </div> <div style="border: 1px solid black; padding: 5px; min-height: 40px;">         Authorized officer           Ting Zhao       </div>
---	--

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/06864

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 435/69.1, 252.2, 254.11, 320.1, 325; 514/44; 530/350; 536/23.5

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 252.2, 254.11, 320.1, 325; 514/44; 530/350; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	FLEURY et al. Uncoupling protein-2: a novel gene linked to obesity and hyperinsulinemia. Nature Genetics. 15 March 1997. Vol. 15, No. 3, pages 269-272, especially the Abstract and Figure 1.	9, 10, 12, 13
X	RAIMBAULT et al. Mus musculus UCP2 mRNA, complete cds. GenBank, Accession Number U69135. 30 October 1996. See the entire document, especially residues 813-827.	9, 12
X ----- Y	HILLIER et al. yf39c01.r1 Homo sapiens cDNA clone 129216 5' similar to SP:UCP_RABIT P14271 MITOCHONDRIAL BROWN FAT UNCOUPLING PROTEIN. EST-STS, Accession Number R11086. 11 April 1995. See the entire document, especially under Source.	12, 13, 15, 16 ----- 1, 2, 4, 177

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 AUGUST 1997

Date of mailing of the international search report

03 SEP 1997

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN LATHROP

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/06864

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHARNOCK-JONES et al. Extension of incomplete cDNAs (ESTs) by biotin/streptavidin-mediated walking using the polymerase chain reaction. Journal of Biotechnology. 30 June 1994. Vol. 35, pages 205-215, see the entire document.	1, 2, 4, 15-177
Y	US 5,453,270 A (BILLS) 26 September 1995, column 2, line 24, through column 3, line 14, and column 3, line 53, through column 4, line 19.	1, 2, 4, 15-17
Y	KOZAK et al. The mitochondrial uncoupling protein gene. The Journal of Biological Chemistry. 05 September 1988. Vol. 263, No. 25, pages 12274-12277. See the entire article.	1, 2, 4, 15-17



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/06864

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 11 and 14  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
Please See Extra Sheet.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-4 and 9-10,12-13,15-17

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

International application No.  
PCT/US97/06864

Claims 11 and 14 are unsearchable to the extent that they require reference to the specified sequences from the Figures. Because applicant has not furnished a computer-readable sequence listing, no meaningful search of the sequences per se can be carried out by this Authority. However, the subject matter of the claims has been searched to the extent possible with reference to the balance of the description.